

RESULTS RECIPIENT
SEATTLE SPERM BANK

Attn: Jeffrey Olliffe 4915 25th Ave NE Ste 204w Seattle, WA 98105-5668

Phone: (206) 588-1484 Fax: (206) 466-4696 NPI: 1306838271 Report Date: 10/05/2020 MALE DONOR 10497

DOB: Ethnicity: Hispanic

Sample Type: EDTA Blood Date of Collection: 09/25/2020 Date Received: 09/26/2020 Date Tested: 10/03/2020 Barcode: 11004512688138 Accession ID: CSLNPJ9A49FQFNQ

Indication: Egg or sperm donor

FEMALE N/A

# Foresight® Carrier Screen

### **NEGATIVE**

### **ABOUT THIS TEST**

The **Myriad Foresight Carrier Screen** utilizes sequencing, maximizing coverage across all DNA regions tested, to help you learn about your chance to have a child with a genetic disease.

#### **RESULTS SUMMARY**

Risk Details	<b>DONOR 10497</b>	Partner
Panel Information	Foresight Carrier Screen Universal Panel Fundamental Plus Panel Fundamental Panel (175 conditions tested)	N/A
All conditions tested A complete list of all conditions tested can be found on page 4.	<ul> <li>□ NEGATIVE</li> <li>No disease-causing mutations were detected.</li> </ul>	N/A

#### **CLINICAL NOTES**

• DONOR is a carrier of one or more extra copies of the alpha globin gene. This alone is not expected to cause clinical signs or symptoms. However, individuals who have one or more extra copies of the alpha globin gene in combination with certain disease-causing HBB variants (beta-plus or beta-zero) may be at risk for a beta thalassemia intermedia phenotype. This phenotype is variable and may result in no symptoms. If not already performed, beta globin gene testing is recommended for a reproductive partner to assess the risk of having a child with beta thalassemia intermedia. Genetic counseling is recommended. See risk calculations at end of report for additional information.

#### **NEXT STEPS**

 If necessary, patients can discuss residual risks with their physician or a genetic counselor.



RESULTS RECIPIENT **SEATTLE SPERM BANK** Attn: Jeffrey Olliffe NPI: 1306838271

Report Date: 10/05/2020

MALE **DONOR 10497** DOB:

Ethnicity: Hispanic Barcode: 11004512688138 FEMALE N/A

## Methods and Limitations

DONOR 10497 [Foresight Carrier Screen]: Sequencing with copy number analysis, spinal muscular atrophy, and analysis of homologous regions (DTS v3.1).

### Sequencing with copy number analysis

High-throughput sequencing and read depth-based copy number analysis are used to analyze the listed exons, as well as selected intergenic and intronic regions, of the genes in the Conditions Tested section of the report. The region of interest (ROI) of the test comprises these regions, in addition to the 20 intronic bases flanking each exon. In a minority of cases where genomic features (e.g., long homopolymers) compromise calling fidelity, the affected intronic bases are not included in the ROI. The ROI is sequenced to high coverage and the sequences are compared to standards and references of normal variation (Genome Reference Consortium Human Build 37 (GRCh37)/hg19). More than 99% of all bases in the ROI are sequenced at greater than the minimum read depth. Mutations may not be detected in areas of lower sequence coverage. Small insertions and deletions may not be as accurately determined as single nucleotide variants. Genes that have closely related pseudogenes may be addressed by a different method. CFTR and DMD testing includes analysis for both large (exon-level) deletions and duplications with an average sensitivity of 99%, while other genes are only analyzed for large deletions with a sensitivity of >75%. However, the sensitivity may be higher for selected founder deletions. The breakpoints of copy number variants and exons affected are estimated from probe positions. Only exons known to be included in the copy number variant are provided in the name. In some cases, the copy number variant may be larger or smaller than indicated. If GJB2 is tested, large upstream deletions involving the genes GJB6 and/or CRYL1 that affect the expression of GJB2 are also analyzed. Mosaicism or somatic variants present at low levels may not be detected. If detected, these may not be reported.

Detection rates are determined by using literature to estimate the fraction of disease alleles, weighted by frequency, that the methodology is unable to detect. Detection rates only account for analytical sensitivity and certain variants that have been previously described in the literature may not be reported if there is insufficient evidence for pathogenicity. Detection rates do not account for the disease-specific rates of de novo mutations.

All variants that are a recognized cause of the disease will be reported. In addition, variants that have not previously been established as a recognized cause of disease may be identified. In these cases, only variants classified as "likely" pathogenic are reported. Likely pathogenic variants are described elsewhere in the report as "likely to have a negative impact on gene function". Likely pathogenic variants are evaluated and classified by assessing the nature of the variant and reviewing reports of allele frequencies in cases and controls, functional studies, variant annotation and effect prediction, and segregation studies. Exon level duplications are assumed to be in tandem and are classified according to their predicted effect on the reading frame. Benign variants, variants of uncertain significance, and variants not directly associated with the intended disease phenotype are not reported. Curation summaries of reported variants are available upon request.

### Spinal muscular atrophy

Targeted copy number analysis is used to determine the copy number of exon 7 of the SMN1 gene relative to other genes. Other mutations may interfere with this analysis. Some individuals with two copies of SMN1 are carriers with two SMN1 genes on one chromosome and a SMN1 deletion on the other chromosome. This is more likely in individuals who have 2 copies of the SMN1 gene and are positive for the g.27134T>G SNP, which affects the reported residual risk; Ashkenazi Jewish or Asian patients with this genotype have a high post-test likelihood of being carriers for SMA and are reported as carriers. The g.27134T>G SNP is only reported in individuals who have 2 copies of SMN1.

### Analysis of homologous regions

A combination of high-throughput sequencing, read depth-based copy number analysis, and targeted genotyping is used to determine the number of functional gene copies and/or the presence of selected loss of function mutations in certain genes that have homology to other regions. The precise breakpoints of large deletions in these genes cannot be determined, but are estimated from copy number analysis. High numbers of pseudogene copies may interfere with this analysis.

If CYP21A2 is tested, patients who have one or more additional copies of the CYP21A2 gene and a loss of function mutation may not actually be a carrier of 21-hydroxylasedeficient congenital adrenal hyperplasia (CAH). Because the true incidence of non-classic CAH is unknown, the residual carrier and reproductive risk numbers on the report are only based on published incidences for classic CAH. However, the published prevalence of non-classic CAH is highest in individuals of Ashkenazi Jewish, Hispanic, Italian, and Yugoslav descent. Therefore, the residual and reproductive risks are likely an underestimate of overall chances for 21-hydroxylase-deficient CAH, especially in the aforementioned populations, as they do not account for non-classic CAH. If HBA1/HBA2 are tested, some individuals with four alpha globin genes may be carriers, with three genes on one chromosome and a deletion on the other chromosome. This and similar, but rare, carrier states, where complementary changes exist in both the gene and a pseudogene, may not be detected by the assay.



MALE
DONOR 10497
DOB:

Ethnicity: Hispanic Barcode: 11004512688138 FEMALE N/A

### Limitations

In an unknown number of cases, nearby genetic variants may interfere with mutation detection. Other possible sources of diagnostic error include sample mix-up, trace contamination, bone marrow transplantation, blood transfusions and technical errors. This test is designed to detect and report germline alterations. While somatic variants present at low levels may be detected, these may not be reported. If more than one variant is detected in a gene, additional studies may be necessary to determine if those variants lie on the same chromosome or different chromosomes. This test is not designed to detect sex chromosome copy number variations. If present, sex chromosome abnormalities may significantly reduce test sensitivity for X-linked conditions. Residual and reproductive risks provided assume a normal karyotype. Risks for individuals with abnormal karyotypes may be different. The test does not fully address all inherited forms of intellectual disability, birth defects and genetic disease. A family history of any of these conditions may warrant additional evaluation. Furthermore, not all mutations will be identified in the genes analyzed and additional testing may be beneficial for some patients. For example, individuals of African, Southeast Asian, and Mediterranean ancestry are at increased risk for being carriers for hemoglobinopathies, which can be identified by CBC and hemoglobin electrophoresis or HPLC (ACOG Practice Bulletin No. 78. Obstet. Gynecol. 2007;109:229-37).

This test was developed and its performance characteristics determined by Myriad Women's Health, Inc. It has not been cleared or approved by the US Food and Drug Administration (FDA). The FDA does not require this test to go through premarket review. This test is used for clinical purposes. It should not be regarded as investigational or for research. This laboratory is certified under the Clinical Laboratory Improvement Amendments of 1988 (CLIA) as qualified to perform high-complexity clinical testing. These results are adjunctive to the ordering physician's evaluation. CLIA Number: #05D1102604.

### Resources

### GENOME CONNECT | http://www.genomeconnect.org

Patients can share their reports via research registries such as Genome Connect, an online research registry working to build the knowledge base about genetics and health. Genome Connect provides patients, physicians, and researchers an opportunity to share genetic information to support the study of the impact of genetic variation on health conditions.

SENIOR LABORATORY DIRECTOR

Jack Ji, PhD, FACMG

Salk

Report content approved by Jack Ji, PhD, FACMG on Oct 5, 2020



MALE DONOR 10497

DOB: Ethnicity: Hispanic

Barcode: 11004512688138

FEMALE N/A

# **Conditions Tested**

11-beta-hydroxylase-deficient Congenital Adrenal Hyperplasia - Gene: CYP11B1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000497:1-9. Detection Rate: Hispanic 94%.

**6-pyruvoyl-tetrahydropterin Synthase Deficiency** - Gene: PTS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_000317:1-6. **Detection Rate:** Hispanic >99%.

ABCC8-related Familial Hyperinsulinism - Gene: ABCC8. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000352:1-39. Detection Rate: Hispanic >99%

Adenosine Deaminase Deficiency - Gene: ADA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000022:1-12. Detection Rate: Hispanic >99%.

Alpha Thalassemia, HBA1/HBA2-related - Genes: HBA1, HBA2. Autosomal Recessive. Analysis of homologous regions. Variants (13): -(alpha)20.5, --BRIT, --MEDI, --MEDII, --SEA, --THAI or --FIL, -alpha3.7, -alpha4.2, HBA1+HBA2 deletion, Hb Constant Spring, anti3.7, anti4.2, del HS-40. Detection Rate: Unknown due to rarity of disease.

Alpha-mannosidosis - Gene: MAN2B1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000528:1-23. Detection Rate: Hispanic >99%.

Alpha-sarcoglycanopathy - Gene: SGCA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000023:1-9. Detection Rate: Hispanic >99%.

Alstrom Syndrome - Gene: ALMS1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_015120:1-23. Detection Rate: Hispanic >99%.

AMT-related Glycine Encephalopathy - Gene: AMT. Autosomal Recessive.

Sequencing with copy number analysis. Exons: NM\_000481:1-9. Detection Rate:

Andermann Syndrome - Gene: SLC12A6. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_133647:1-25. Detection Rate: Hispanic >99%. Argininemia - Gene: ARG1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000045:1-8. Detection Rate: Hispanic 97%.

**Argininosuccinic Aciduria** - **Gene**: ASL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_001024943:1-16. **Detection Rate:** Hispanic >99%.

**Aspartylglucosaminuria** - Gene: AGA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000027:1-9. Detection Rate: Hispanic >99%.

Ataxia with Vitamin E Deficiency - Gene: TTPA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000370:1-5. Detection Rate: Hispanic >99%. Ataxia-telangiectasia - Gene: ATM. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000051:2-63. Detection Rate: Hispanic 97%.

**ATP7A-related Disorders** - **Gene:** ATP7A. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM\_000052:2-23. **Detection Rate:** Hispanic 92%.

Autoimmune Polyglandular Syndrome Type 1 - Gene: AIRE. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000383:1-14. Detection Rate: Hispanic >99%.

Autosomal Recessive Osteopetrosis Type 1 - Gene: TCIRG1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_006019:2-20. Detection Rate: Hispanic >99%.

Autosomal Recessive Polycystic Kidney Disease, PKHD1-related - Gene: PKHD1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_138694 2-67. Detection Rate: Hispanic >99%.

Autosomal Recessive Spastic Ataxia of Charlevoix-Saguenay - Gene: SACS. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_014363 2-10. Detection Rate: Hispanic 99%.

Bardet-Biedl Syndrome, BBS1-related - Gene: BBS1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_024649:1-17. Detection Rate: Hispanic >99%.

Bardet-Biedl Syndrome, BBS10-related - Gene: BBS10. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_024685:1-2. Detection Rate: Hispanic >99%.

**Bardet-Biedl Syndrome, BBS12-related** - Gene: BBS12. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM\_152618:2. **Detection Rate:** Hispanic >99%.

Bardet-Biedl Syndrome, BBS2-related - Gene: BBS2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_031885:1-17. Detection Rate: Hispanic >99%.

BCS1L-related Disorders - Gene: BCS1L. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_004328:3-9. Detection Rate: Hispanic >99%. Beta-sarcoglycanopathy - Gene: SGCB. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000232:1-6. Detection Rate: Hispanic >99%. Biotinidase Deficiency - Gene: BTD. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000060:1-4. Detection Rate: Hispanic >99%. Bloom Syndrome - Gene: BLM. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000057:2-22. Detection Rate: Hispanic >99%. Calpainopathy - Gene: CAPN3. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000070:1-24. Detection Rate: Hispanic >99%. Canavan Disease - Gene: ASPA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000049:1-6. Detection Rate: Hispanic 98%. Carbamoylphosphate Synthetase I Deficiency - Gene: CPS1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_0001875:1-38. Detection Rate:

Carnitine Palmitoyltransferase IA Deficiency - Gene: CPT1A. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_001876:2-19. Detection Rate: Hispanic >99%.

Carnitine Palmitoyltransferase II Deficiency - Gene: CPT2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000098:1-5. Detection Rate: Hispanic >99%.

Cartilage-hair Hypoplasia - Gene: RMRP. Autosomal Recessive. Sequencing with copy number analysis. Exon: NR\_003051:1. Detection Rate: Hispanic >99%. Cerebrotendinous Xanthomatosis - Gene: CYP27A1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000784:1-9. Detection Rate: Hispanic >99%.

Citrullinemia Type 1 - Gene: ASS1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000050:3-16. Detection Rate: Hispanic >99%. CLN3-related Neuronal Ceroid Lipofuscinosis - Gene: CLN3. Autosomal Recessive.

Sequencing with copy number analysis. Exons: NM\_001042432 2-16. Detection Rate: Hispanic >99%.

**CLN5-related Neuronal Ceroid Lipofuscinosis** - Gene: CLN5. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_006493:1-4. **Detection Rate:** Hispanic >99%.

**CLN6-related Neuronal Ceroid Lipofuscinosis** - Gene: CLN6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_017882:1-7. **Detection Rate:** Hispanic >99%.

CLN8-related Neuronal Ceroid Lipofuscinosis - Gene: CLN8. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_018941:2-3. Detection Rate: Hispanic >99%.

Cohen Syndrome - Gene: VPS13B. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_017890:2-62. Detection Rate: Hispanic 97%. COL4A3-related Alport Syndrome - Gene: COL4A3. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000091:1-52. Detection Rate: Hispanic 97%.

**COL4A4**-related Alport Syndrome - Gene: COL4A4. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000092:2-48. Detection Rate: Hispanic 98%.

Combined Pituitary Hormone Deficiency, PROP1-related - Gene: PROP1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_006261:1-3. Detection Rate: Hispanic >99%.



MALE DONOR 10497

DOB: Ethnicity: Hispanic

Barcode: 11004512688138

FEMALE N/A

Congenital Adrenal Hyperplasia, CYP21A2-related - Gene: CYP21A2. Autosomal Recessive. Analysis of homologous regions. Variants (13): CYP21A2 deletion, CYP21A2 duplication, CYP21A2 triplication, G111Vfs\*21, I173N, L308Ffs\*6, P31L, Q319\*, Q319\*+CYP21A2dup, R357W, V281L, [I237N;V238E;M240K], c.293-13C>G. Detection Rate: Hispanic 95%.

Congenital Disorder of Glycosylation Type Ia - Gene: PMM2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000303:1-8. Detection Rate: Hispanic >99%.

Congenital Disorder of Glycosylation Type Ic - Gene: ALG6. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_013339:2-15. Detection Rate: Hispanic >99%.

Congenital Disorder of Glycosylation, MPI-related - Gene: MPI. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_002435:1-8. Detection Rate: Hispanic >99%.

Costeff Optic Atrophy Syndrome - Gene: OPA3. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_025136:1-2. Detection Rate: Hispanic >99%. Cystic Fibrosis - Gene: CFTR. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000492:1-27. IVS8-5T allele analysis is only reported in the presence of the R117H mutation. Detection Rate: Hispanic >99%.

**Cystinosis** - **Gene**: CTNS. Autosomal Recessive. Sequencing with copy number analysis. **Exons**: NM\_004937:3-12. **Detection Rate**: Hispanic >99%.

**D-bifunctional Protein Deficiency** - Gene: HSD17B4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_000414:1-24. **Detection Rate:** Hispanic 98%.

**Delta-sarcoglycanopathy** - **Gene**: SGCD. Autosomal Recessive. Sequencing with copy number analysis. **Exons**: NM\_000337:2-9. **Detection Rate**: Hispanic 99%.

**Dihydrolipoamide Dehydrogenase Deficiency** - Gene: DLD. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000108:1-14. Detection Rate: Hispanic >99%.

**Dysferlinopathy** - **Gene**: DYSF. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_003494:1-55. **Detection Rate**: Hispanic 98%.

Dystrophinopathy (Including Duchenne/Becker Muscular Dystrophy) - Gene: DMD. X-linked Recessive. Sequencing with copy number analysis. Exons: NM\_004006:1-79. Detection Rate: Hispanic >99%.

ERCC6-related Disorders - Gene: ERCC6. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000124:2-21. Detection Rate: Hispanic 99%. ERCC8-related Disorders - Gene: ERCC8. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000082:1-12. Detection Rate: Hispanic 95%.

**EVC-related Ellis-van Creveld Syndrome** - Gene: EVC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_153717:1-21. **Detection Rate:** Hispanic 96%.

**EVC2-related Ellis-van Creveld Syndrome** - Gene: EVC2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_147127:1-22. **Detection Rate:** Hispanic >99%.

**Fabry Disease** - Gene: GLA. X-linked Recessive. Sequencing with copy number analysis. Exons: NM\_000169:1-7. Detection Rate: Hispanic 98%.

Familial Dysautonomia - Gene: IKBKAP. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_003640:2-37. Detection Rate: Hispanic >99%. Familial Mediterranean Fever - Gene: MEFV. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000243:1-10. Detection Rate: Hispanic >00%

Fanconi Anemia Complementation Group A - Gene: FANCA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000135:1-43. Detection Rate: Hispanic 92%.

Fanconi Anemia, FANCC-related - Gene: FANCC. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000136:2-15. Detection Rate: Hispanic >99%.

**FKRP-related Disorders** - Gene: FKRP. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM\_024301:4. **Detection Rate:** Hispanic >99%.

**FKTN-related Disorders** - Gene: FKTN. Autosomal Recessive. Sequencing with copy number analysis. **Exons**: NM\_001079802:3-11. **Detection Rate**: Hispanic >99%.

Free Sialic Acid Storage Disorders - Gene: SLC17A5. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_012434:1-11. Detection Rate: Hispanic 98%.

Galactokinase Deficiency - Gene: GALK1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000154:1-8. Detection Rate: Hispanic >99%.

**Galactosemia** - Gene: GALT. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000155:1-11. Detection Rate: Hispanic >99%.

Gamma-sarcoglycanopathy - Gene: SGCG. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000231:2-8. Detection Rate: Hispanic 88%. Gaucher Disease - Gene: GBA. Autosomal Recessive. Analysis of homologous regions. Variants (10): D409V, D448H, IVS2+1G>A, L444P, N370S, R463C, R463H, R496H, V394L, p.L29Afs\*18. Detection Rate: Hispanic 60%.

GJB2-related DFNB1 Nonsyndromic Hearing Loss and Deafness - Gene: GJB2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_004004:1-2. Detection Rate: Hispanic >99%.

GLB1-related Disorders - Gene: GLB1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000404:1-16. Detection Rate: Hispanic >99%.
GLDC-related Glycine Encephalopathy - Gene: GLDC. Autosomal Recessive.
Sequencing with copy number analysis. Exons: NM\_000170:1-25. Detection Rate: Hispanic 94%.

**Glutaric Acidemia, GCDH-related** - Gene: GCDH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_000159:2-12. **Detection Rate:** Hispanic >99%.

**Glycogen Storage Disease Type Ia** - **Gene**: G6PC. Autosomal Recessive. Sequencing with copy number analysis. **Exons**: NM\_000151:1-5. **Detection Rate**: Hispanic >99%.

**Glycogen Storage Disease Type Ib - Gene:** SLC37A4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_001164277 3-11. **Detection Rate:** Hispanic >99%.

**Glycogen Storage Disease Type III** - Gene: AGL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_000642:2-34. **Detection Rate:** Hispanic >99%

**GNE Myopathy** - **Gene:** GNE. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_001128227:1-12. **Detection Rate:** Hispanic >99%.

**GNPTAB-related Disorders** - Gene: GNPTAB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_024312:1-21. **Detection Rate:** Hispanic >00%

HADHA-related Disorders - Gene: HADHA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000182:1-20. Detection Rate: Hispanic >99%. Hb Beta Chain-related Hemoglobinopathy (Including Beta Thalassemia and Sickle Cell Disease) - Gene: HBB. Autosomal Recessive. Sequencing with copy number

analysis. Exons: NM\_000518:1-3. Detection Rate: Hispanic >99%. Hereditary Fructose Intolerance - Gene: ALDOB. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000035:2-9. Detection Rate: Hispanic >99%.

Herlitz Junctional Epidermolysis Bullosa, LAMB3-related - Gene: LAMB3.

Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000228 2-23. Detection Rate: Hispanic >99%.

Hexosaminidase A Deficiency (Including Tay-Sachs Disease) - Gene: HEXA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000520:1-14. Detection Rate: Hispanic >99%.

HMG-CoA Lyase Deficiency - Gene: HMGCL. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000191:1-9. Detection Rate: Hispanic 98%. Holocarboxylase Synthetase Deficiency - Gene: HLCS. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000411:4-12. Detection Rate: Hispanic >99%.

Homocystinuria, CBS-related - Gene: CBS. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000071:3-17. Detection Rate: Hispanic >99%. Hydrolethalus Syndrome - Gene: HYLS1. Autosomal Recessive. Sequencing with copy number analysis. Exon: NM\_145014:4. Detection Rate: Hispanic >99%. Hypophosphatasia - Gene: ALPL. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000478:2-12. Detection Rate: Hispanic >99%. Isovaleric Acidemia - Gene: IVD. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_002225:1-12. Detection Rate: Hispanic >99%.



MALE DONOR 10497

DOB: Ethnicity: Hispanic

Barcode: 11004512688138

FEMALE N/A

Joubert Syndrome 2 - Gene: TMEM216. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_001173990:1-5. Detection Rate: Hispanic >99%. Junctional Epidermolysis Bullosa, LAMA3-related - Gene: LAMA3. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000227:1-38. Detection Rate: Hispanic >99%.

Junctional Epidermolysis Bullosa, LAMC2-related - Gene: LAMC2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_005562:1-23. Detection Rate: Hispanic >99%.

KCNJ11-related Familial Hyperinsulinism - Gene: KCNJ11. Autosomal Recessive. Sequencing with copy number analysis. Exon: NM\_000525:1. Detection Rate: Hispanic >99%.

Krabbe Disease - Gene: GALC. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000153:1-17. Detection Rate: Hispanic >99%. LAMA2-related Muscular Dystrophy - Gene: LAMA2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000426:1-65. Detection Rate: Hispanic >99%.

**Leigh Syndrome, French-Canadian Type** - Gene: LRPPRC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_133259:1-38. **Detection Rate:** Hispanic >99%.

**Lipoid Congenital Adrenal Hyperplasia** - Gene: STAR. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000349:1-7. Detection Rate: Hispanic >99%.

Lysosomal Acid Lipase Deficiency - Gene: LIPA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000235:2-10. Detection Rate: Hispanic >99%.

Maple Syrup Urine Disease Type Ia - Gene: BCKDHA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000709:1-9. Detection Rate: Hispanic >99%.

Maple Syrup Urine Disease Type Ib - Gene: BCKDHB. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_183050:1-10. Detection Rate: Hispanic >99%.

Maple Syrup Urine Disease Type II - Gene: DBT. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_001918:1-11. Detection Rate: Hispanic 96%. Medium Chain Acyl-CoA Dehydrogenase Deficiency - Gene: ACADM. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000016:1-12. Detection Rate: Hispanic >99%.

Megalencephalic Leukoencephalopathy with Subcortical Cysts - Gene: MLC1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_015166 2-12. Detection Rate: Hispanic >99%.

Metachromatic Leukodystrophy - Gene: ARSA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000487:1-8. Detection Rate: Hispanic >99%. Methylmalonic Acidemia, cblA Type - Gene: MMAA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_172250:2-7. Detection Rate: Hispanic >99%.

Methylmalonic Acidemia, cblB Type - Gene: MMAB. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_052845:1-9. Detection Rate: Hispanic >99%.

Methylmalonic Aciduria and Homocystinuria, cblC Type - Gene: MMACHC. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_015506:1-4. Detection Rate: Hispanic >99%.

MKS1-related Disorders - Gene: MKS1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_017777:1-18. Detection Rate: Hispanic >99%. Mucolipidosis III Gamma - Gene: GNPTG. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_032520:1-11. Detection Rate: Hispanic >99%. Mucolipidosis IV - Gene: MCOLN1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_020533:1-14. Detection Rate: Hispanic >99%.

**Mucopolysaccharidosis Type I** - Gene: IDUA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_000203:1-14. **Detection Rate:** Hispanic >99%.

Mucopolysaccharidosis Type II - Gene: IDS. X-linked Recessive. Sequencing with copy number analysis. Exons: NM\_000202:1-9. Detection Rate: Hispanic 88%.

Mucopolysaccharidosis Type IIIA - Gene: SGSH. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000199:1-8. Detection Rate: Hispanic >99%.

**Mucopolysaccharidosis Type IIIB** - Gene: NAGLU. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000263:1-6. **Detection Rate:** Hispanic >99%.

**Mucopolysaccharidosis Type IIIC** - Gene: HGSNAT. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_152419:1-18. Detection Rate: Hispanic >99%.

MUT-related Methylmalonic Acidemia - Gene: MUT. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000255:2-13. Detection Rate: Hispanic >99%.

MYO7A-related Disorders - Gene: MYO7A. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000260:2-49. Detection Rate: Hispanic >99%. NEB-related Nemaline Myopathy - Gene: NEB. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_001271208:3-80,117-183. Detection Rate: Hispanic 92%.

**Nephrotic Syndrome, NPHS1-related** - Gene: NPHS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_004646:1-29. **Detection Rate:** Hispanic >99%.

**Nephrotic Syndrome, NPHS2-related** - Gene: NPHS2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_014625:1-8. **Detection Rate:** Hispanic >99%.

Niemann-Pick Disease Type C1 - Gene: NPC1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000271:1-25. Detection Rate: Hispanic >99%.

Niemann-Pick Disease Type C2 - Gene: NPC2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_006432:1-5. Detection Rate: Hispanic >99%. Niemann-Pick Disease, SMPD1-related - Gene: SMPD1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000543:1-6. Detection Rate: Hispanic >99%.

Nijmegen Breakage Syndrome - Gene: NBN. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_002485:1-16. Detection Rate: Hispanic >99%.

Ornithine Transcarbamylase Deficiency - Gene: OTC. X-linked Recessive. Sequencing with copy number analysis. Exons: NM\_000531:1-10. Detection Rate: Hispanic 97%.

PCCA-related Propionic Acidemia - Gene: PCCA. Autosomal Recessive.
Sequencing with copy number analysis. Exons: NM\_000282:1-24. Detection Rate: Hispanic 95%.

PCCB-related Propionic Acidemia - Gene: PCCB. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000532:1-15. Detection Rate: Hispanic >99%.

PCDH15-related Disorders - Gene: PCDH15. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_033056:2-33. Detection Rate: Hispanic 93%. Pendred Syndrome - Gene: SLC26A4. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000441:2-21. Detection Rate: Hispanic >99%.

**Peroxisome Biogenesis Disorder Type 1** - Gene: PEX1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_000466:1-24. **Detection Rate:** Hispanic >99%.

**Peroxisome Biogenesis Disorder Type 3** - Gene: PEX12. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000286:1-3. **Detection Rate:** Hispanic >99%.

**Peroxisome Biogenesis Disorder Type 4** - Gene: PEX6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_000287:1-17. **Detection Rate:** Hispanic 97%.

**Peroxisome Biogenesis Disorder Type 5** - Gene: PEX2. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM\_000318:4. **Detection Rate:** Hispanic >99%.

**Peroxisome Biogenesis Disorder Type 6** - Gene: PEX10. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_153818:1-6. **Detection Rate:** Hispanic >99%.

Phenylalanine Hydroxylase Deficiency - Gene: PAH. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000277:1-13. Detection Rate: Hispanic >99%.



MALE DONOR 10497

DOB: Ethnicity: Hispanic

Barcode: 11004512688138

FEMALE N/A

POMGNT-related Disorders - Gene: POMGNT1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_017739:2-22. Detection Rate: Hispanic 96%. Pompe Disease - Gene: GAA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000152:2-20. Detection Rate: Hispanic 98%.

**PPT1-related Neuronal Ceroid Lipofuscinosis** - Gene: PPT1. Autosomal Recessive. Sequencing with copy number analysis. **Exons**: NM\_000310:1-9. **Detection Rate**: Hispanic >99%.

Primary Carnitine Deficiency - Gene: SLC22A5. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_003060:1-10. Detection Rate: Hispanic >99%

**Primary Hyperoxaluria Type 1** - **Gene:** AGXT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_000030:1-11. **Detection Rate:** Hispanic >99%.

Primary Hyperoxaluria Type 2 - Gene: GRHPR. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_012203:1-9. Detection Rate: Hispanic >99%. Primary Hyperoxaluria Type 3 - Gene: HOGA1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_138413:1-7. Detection Rate: Hispanic >99%. Pycnodysostosis - Gene: CTSK. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000396:2-8. Detection Rate: Hispanic >99%. Pyruvate Carboxylase Deficiency - Gene: PC. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000920:3-22. Detection Rate: Hispanic >99%.

Rhizomelic Chondrodysplasia Punctata Type 1 - Gene: PEX7. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000288:1-10. Detection Rate: Hispanic >99%.

RTEL1-related Disorders - Gene: RTEL1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_032957:2-35. Detection Rate: Hispanic >99%. Sandhoff Disease - Gene: HEXB. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000521:1-14. Detection Rate: Hispanic 99%. Short-chain Acyl-CoA Dehydrogenase Deficiency - Gene: ACADS. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000017:1-10. Detection Rate: Hispanic >99%.

Sjogren-Larsson Syndrome - Gene: ALDH3A2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000382:1-10. Detection Rate: Hispanic 96%. SLC26A2-related Disorders - Gene: SLC26A2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000112:2-3. Detection Rate: Hispanic >99%. Smith-Lemli-Opitz Syndrome - Gene: DHCR7. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_001360:3-9. Detection Rate: Hispanic >99%. Spastic Paraplegia Type 15 - Gene: ZFYVE26. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_015346:2-42. Detection Rate: Hispanic >99%.

Spinal Muscular Atrophy - Gene: SMN1. Autosomal Recessive. Spinal muscular atrophy. Variant (1): SMN1 copy number. Detection Rate: Hispanic 91%. Spondylothoracic Dysostosis - Gene: MESP2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_001039958:1-2. Detection Rate: Hispanic >99%.

**TGM1-related Autosomal Recessive Congenital Ichthyosis** - Gene: TGM1. Autosomal Recessive. Sequencing with copy number analysis. Exons:

NM\_000359 2-15. Detection Rate: Hispanic >99%.

**TPP1-related Neuronal Ceroid Lipofuscinosis** - Gene: TPP1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_000391:1-13. **Detection Rate:** Hispanic >99%.

**Tyrosine Hydroxylase Deficiency** - **Gene:** TH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM\_199292:1-14. **Detection Rate:** Hispanic >99%.

Tyrosinemia Type I - Gene: FAH. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000137:1-14. Detection Rate: Hispanic >99%.

Tyrosinemia Type II - Gene: TAT. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000353:2-12. Detection Rate: Hispanic >99%.

USH1C-related Disorders - Gene: USH1C. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_005709:1-21. Detection Rate: Hispanic >99%.

USH2A-related Disorders - Gene: USH2A. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_206933:2-72. Detection Rate: Hispanic 94%.

Usher Syndrome Type 3 - Gene: CLRN1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_174878:1-3. Detection Rate: Hispanic >99%.

Very-long-chain Acyl-CoA Dehydrogenase Deficiency - Gene: ACADVL. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000018:1-20.

Detection Rate: Hispanic >99%.

Wilson Disease - Gene: ATP7B. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000053:1-21. Detection Rate: Hispanic >99%.

X-linked Adrenoleukodystrophy - Gene: ABCD1. X-linked Recessive. Sequencing with copy number analysis. Exons: NM\_000033:1-6. Detection Rate: Hispanic 77%.

X-linked Alport Syndrome - Gene: COL4A5. X-linked Recessive. Sequencing with copy number analysis. Exons: NM\_000495:1-51. Detection Rate: Hispanic 95%.

X-linked Congenital Adrenal Hypoplasia - Gene: NR0B1. X-linked Recessive.

Sequencing with copy number analysis. Exons: NM\_000475:1-2. Detection Rate: Hispanic 99%.

X-linked Juvenile Retinoschisis - Gene: RS1. X-linked Recessive. Sequencing with copy number analysis. Exons: NM\_000330:1-6. Detection Rate: Hispanic 98%.
X-linked Myotubular Myopathy - Gene: MTM1. X-linked Recessive. Sequencing with copy number analysis. Exons: NM\_000252:2-15. Detection Rate: Hispanic 98%.
X-linked Severe Combined Immunodeficiency - Gene: IL2RG. X-linked Recessive. Sequencing with copy number analysis. Exons: NM\_000206:1-8. Detection Rate: Hispanic >99%.

Xeroderma Pigmentosum Group A - Gene: XPA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000380:1-6. Detection Rate: Hispanic >99%. Xeroderma Pigmentosum Group C - Gene: XPC. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_004628:1-16. Detection Rate: Hispanic 97%.



MALE **DONOR 10497** DOB:

Ethnicity: Hispanic

Barcode: 11004512688138

**FEMALE** N/A

## Risk Calculations

Below are the risk calculations for all conditions tested. Since negative results do not completely rule out the possibility of being a carrier, the residual risk represents the patient's post-test likelihood of being a carrier and the reproductive risk represents the likelihood the patient's future children could inherit each disease. These risks are inherent to all carrier screening tests, may vary by ethnicity, are predicated on a negative family history and are present even after a negative test result. Inaccurate reporting of ethnicity may cause errors in risk calculation. The reproductive risk presented is based on a hypothetical pairing with a partner of the same ethnic group.

Disease	DONOR 10497 Residual Risk	Reproductive Risk
11-beta-hydroxylase-deficient Congenital Adrenal Hyperplasia	1 in 3,300	< 1 in 1,000,000
6-pyruvoyl-tetrahydropterin Synthase Deficiency	< 1 in 50,000	< 1 in 1,000,000
ABCC8-related Familial Hyperinsulinism	1 in 17,000	< 1 in 1,000,000
Adenosine Deaminase Deficiency	1 in 39,000	< 1 in 1,000,000
Alpha Thalassemia, HBA1/HBA2-related	Alpha globin status: aa/aaa.	Not calculated
Alpha-mannosidosis	1 in 35,000	< 1 in 1,000,000
Alpha-sarcoglycanopathy	1 in 34,000	< 1 in 1,000,000
Alstrom Syndrome	< 1 in 50,000	< 1 in 1,000,000
AMT-related Glycine Encephalopathy	1 in 22,000	< 1 in 1,000,000
Andermann Syndrome	< 1 in 50,000	< 1 in 1,000,000
Argininemia	< 1 in 17,000	< 1 in 1,000,000
Argininosuccinic Aciduria	1 in 29,000	< 1 in 1,000,000
Aspartylglucosaminuria	< 1 in 50,000	< 1 in 1,000,000
Ataxia with Vitamin E Deficiency		
·	< 1 in 50,000	< 1 in 1,000,000
Ataxia-telangiectasia	1 in 3,700	< 1 in 1,000,000
ATP7A-related Disorders	< 1 in 1,000,000	1 in 600,000
Autoimmune Polyglandular Syndrome Type 1	1 in 18,000	< 1 in 1,000,000
Autosomal Recessive Osteopetrosis Type 1	1 in 35,000	< 1 in 1,000,000
Autosomal Recessive Polycystic Kidney Disease, PKHD1-related	1 in 8,100	< 1 in 1,000,000
Autosomal Recessive Spastic Ataxia of Charlevoix-Saguenay	< 1 in 44,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS1-related	1 in 39,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS10-related	1 in 42,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS12-related	< 1 in 50,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS2-related	< 1 in 50,000	< 1 in 1,000,000
BCS1L-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Beta-sarcoglycanopathy	1 in 39,000	< 1 in 1,000,000
Biotinidase Deficiency	1 in 17,000	< 1 in 1,000,000
Bloom Syndrome	< 1 in 50,000	< 1 in 1,000,000
Calpainopathy	1 in 13,000	< 1 in 1,000,000
Canavan Disease	1 in 9,700	< 1 in 1,000,000
Carbamoylphosphate Synthetase I Deficiency	< 1 in 57,000	< 1 in 1,000,000
Carnitine Palmitoyltransferase IA Deficiency	< 1 in 50,000	< 1 in 1,000,000
Carnitine Palmitoyltransferase II Deficiency	1 in 18,000	< 1 in 1,000,000
Cartilage-hair Hypoplasia	< 1 in 50,000	< 1 in 1,000,000
Cerebrotendinous Xanthomatosis	1 in 11,000	< 1 in 1,000,000
	1 in 12,000	< 1 in 1,000,000
Citrullinemia Type 1		
CLN3-related Neuronal Ceroid Lipofuscinosis	1 in 13,000	< 1 in 1,000,000
CLN5-related Neuronal Ceroid Lipofuscinosis	< 1 in 50,000	< 1 in 1,000,000
CLN6-related Neuronal Ceroid Lipofuscinosis	< 1 in 50,000	< 1 in 1,000,000
CLN8-related Neuronal Ceroid Lipofuscinosis	< 1 in 50,000	< 1 in 1,000,000
Cohen Syndrome	< 1 in 15,000	< 1 in 1,000,000
COL4A3-related Alport Syndrome	1 in 11,000	< 1 in 1,000,000
COL4A4-related Alport Syndrome	1 in 21,000	< 1 in 1,000,000
Combined Pituitary Hormone Deficiency, PROP1-related	1 in 6,100	< 1 in 1,000,000
Congenital Adrenal Hyperplasia, CYP21A2-related		
Congenital Disorder of Glycosylation Type Ia	1 in 1,200	1 in 290,000
Congenital Disorder of Glycosylation Type Ic		1 in 290,000 < 1 in 1,000,000
Congenital Disorder of Chycosylation Type IC	1 in 1,200	
Congenital Disorder of Glycosylation Type IC	1 in 1,200 1 in 16,000	< 1 in 1,000,000
	1 in 1,200 1 in 16,000 < 1 in 50,000	< 1 in 1,000,000 < 1 in 1,000,000
Congenital Disorder of Glycosylation, MPI-related	1 in 1,200 1 in 16,000 < 1 in 50,000 < 1 in 50,000	< 1 in 1,000,000 < 1 in 1,000,000 < 1 in 1,000,000



MALE DONOR 10497

DOB: Ethnicity: Hispanic

Barcode: 11004512688138

FEMALE N/A

Disease	DONOR 10497 Residual Risk	Reproductive Risk
		•
D-bifunctional Protein Deficiency	1 in 9,000	< 1 in 1,000,000
Delta-sarcoglycanopathy	< 1 in 40,000	< 1 in 1,000,000
Dihydrolipoamide Dehydrogenase Deficiency	< 1 in 50,000	< 1 in 1,000,000
Dysferlinopathy	1 in 11,000	< 1 in 1,000,000
Dystrophinopathy (Including Duchenne/Becker Muscular Dystrophy)	Not calculated	Not calculated
ERCC6-related Disorders	1 in 19,000	< 1 in 1,000,000
ERCC8-related Disorders	1 in 7,300	< 1 in 1,000,000
EVC-related Ellis-van Creveld Syndrome	1 in 7,500	< 1 in 1,000,000
EVC2-related Ellis-van Creveld Syndrome	< 1 in 50,000	< 1 in 1,000,000
Fabry Disease	< 1 in 1,000,000	1 in 80,000
Familial Dysautonomia	< 1 in 50,000	< 1 in 1,000,000
Familial Mediterranean Fever	< 1 in 50,000	< 1 in 1,000,000
Fanconi Anemia Complementation Group A	1 in 2,900	< 1 in 1,000,000
Fanconi Anemia, FANCC-related	< 1 in 50,000	< 1 in 1,000,000
FKRP-related Disorders	1 in 19,000	< 1 in 1,000,000
FKTN-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Free Sialic Acid Storage Disorders	< 1 in 30,000	< 1 in 1,000,000
Galactokinase Deficiency	1 in 35,000	< 1 in 1,000,000
Galactosemia	1 in 11,000	< 1 in 1,000,000
Gamma-sarcoglycanopathy	1 in 3,000	< 1 in 1,000,000
Gaucher Disease	1 in 310	1 in 150,000
GJB2-related DFNB1 Nonsyndromic Hearing Loss and Deafness	1 in 3,200	1 in 410,000
GLB1-related Disorders	1 in 19,000	< 1 in 1,000,000
GLDC-related Glycine Encephalopathy	1 in 2,800	< 1 in 1,000,000
Glutaric Acidemia, GCDH-related	1 in 16,000	< 1 in 1,000,000
Glycogen Storage Disease Type Ia	1 in 18,000	< 1 in 1,000,000
Glycogen Storage Disease Type Ib	1 in 35,000	< 1 in 1,000,000
Glycogen Storage Disease Type III	1 in 16,000	< 1 in 1,000,000
GNE Myopathy	< 1 in 50,000	< 1 in 1,000,000
GNPTAB-related Disorders	1 in 32,000	< 1 in 1,000,000
HADHA-related Disorders	1 in 25,000	< 1 in 1,000,000
Hb Beta Chain-related Hemoglobinopathy (Including Beta Thalassemia and Sickle Ce Disease)	II 1 in 2,400	1 in 240,000
Hereditary Fructose Intolerance	1 in 7,900	< 1 in 1,000,000
Herlitz Junctional Epidermolysis Bullosa, LAMB3-related	< 1 in 50,000	< 1 in 1,000,000
Hexosaminidase A Deficiency (Including Tay-Sachs Disease)	1 in 30,000	< 1 in 1,000,000
HMG-CoA Lyase Deficiency	< 1 in 33,000	< 1 in 1,000,000
Holocarboxylase Synthetase Deficiency	1 in 15,000	< 1 in 1,000,000
Homocystinuria, CBS-related	1 in 10,000	< 1 in 1,000,000
Hydrolethalus Syndrome	< 1 in 50,000	< 1 in 1,000,000
Hypophosphatasia	1 in 22,000	< 1 in 1,000,000
Isovaleric Acidemia	1 in 26,000	< 1 in 1,000,000
Joubert Syndrome 2	< 1 in 50,000	< 1 in 1,000,000
Junctional Epidermolysis Bullosa, LAMA3-related	< 1 in 50,000	< 1 in 1,000,000
Junctional Epidermolysis Bullosa, LAMC2-related	< 1 in 50,000	< 1 in 1,000,000
KCNJ11-related Familial Hyperinsulinism	< 1 in 50,000	< 1 in 1,000,000
Krabbe Disease	1 in 17,000	< 1 in 1,000,000
LAMA2-related Muscular Dystrophy	1 in 17,000	< 1 in 1,000,000
	1 111 17,000	× 1 III 1,000,000
· · · ·	< 1 in 50 000	< 1 in 1 000 000
Leigh Syndrome, French-Canadian Type	< 1 in 50,000	< 1 in 1,000,000
Leigh Syndrome, French-Canadian Type Lipoid Congenital Adrenal Hyperplasia	< 1 in 50,000	< 1 in 1,000,000
Leigh Syndrome, French-Canadian Type Lipoid Congenital Adrenal Hyperplasia Lysosomal Acid Lipase Deficiency	< 1 in 50,000 1 in 18,000	< 1 in 1,000,000 < 1 in 1,000,000
Leigh Syndrome, French-Canadian Type Lipoid Congenital Adrenal Hyperplasia Lysosomal Acid Lipase Deficiency Maple Syrup Urine Disease Type Ia	< 1 in 50,000 1 in 18,000 1 in 14,000	< 1 in 1,000,000 < 1 in 1,000,000 < 1 in 1,000,000
Leigh Syndrome, French-Canadian Type Lipoid Congenital Adrenal Hyperplasia Lysosomal Acid Lipase Deficiency Maple Syrup Urine Disease Type Ia Maple Syrup Urine Disease Type Ib	< 1 in 50,000 1 in 18,000 1 in 14,000 1 in 36,000	< 1 in 1,000,000 < 1 in 1,000,000 < 1 in 1,000,000 < 1 in 1,000,000
Leigh Syndrome, French-Canadian Type Lipoid Congenital Adrenal Hyperplasia Lysosomal Acid Lipase Deficiency Maple Syrup Urine Disease Type Ia Maple Syrup Urine Disease Type Ib Maple Syrup Urine Disease Type II	< 1 in 50,000 1 in 18,000 1 in 14,000 1 in 36,000 1 in 13,000	< 1 in 1,000,000 < 1 in 1,000,000 < 1 in 1,000,000 < 1 in 1,000,000 < 1 in 1,000,000
Leigh Syndrome, French-Canadian Type Lipoid Congenital Adrenal Hyperplasia Lysosomal Acid Lipase Deficiency Maple Syrup Urine Disease Type Ia Maple Syrup Urine Disease Type Ib Maple Syrup Urine Disease Type II Medium Chain Acyl-CoA Dehydrogenase Deficiency	< 1 in 50,000 1 in 18,000 1 in 14,000 1 in 36,000 1 in 13,000 1 in 6,000	< 1 in 1,000,000 < 1 in 1,000,000
Leigh Syndrome, French-Canadian Type Lipoid Congenital Adrenal Hyperplasia Lysosomal Acid Lipase Deficiency Maple Syrup Urine Disease Type Ia Maple Syrup Urine Disease Type Ib Maple Syrup Urine Disease Type II Medium Chain Acyl-CoA Dehydrogenase Deficiency Megalencephalic Leukoencephalopathy with Subcortical Cysts	< 1 in 50,000 1 in 18,000 1 in 14,000 1 in 36,000 1 in 13,000 1 in 6,000 < 1 in 50,000	< 1 in 1,000,000 < 1 in 1,000,000
Leigh Syndrome, French-Canadian Type Lipoid Congenital Adrenal Hyperplasia Lysosomal Acid Lipase Deficiency Maple Syrup Urine Disease Type Ia Maple Syrup Urine Disease Type Ib Maple Syrup Urine Disease Type II Medium Chain Acyl-CoA Dehydrogenase Deficiency Megalencephalic Leukoencephalopathy with Subcortical Cysts Metachromatic Leukodystrophy	< 1 in 50,000 1 in 18,000 1 in 14,000 1 in 36,000 1 in 13,000 1 in 6,000 < 1 in 50,000 1 in 16,000	<pre>&lt; 1 in 1,000,000 &lt; 1 in 1,000,000</pre>
Leigh Syndrome, French-Canadian Type Lipoid Congenital Adrenal Hyperplasia Lysosomal Acid Lipase Deficiency Maple Syrup Urine Disease Type Ia Maple Syrup Urine Disease Type Ib Maple Syrup Urine Disease Type II Medium Chain Acyl-CoA Dehydrogenase Deficiency Megalencephalic Leukoencephalopathy with Subcortical Cysts Metachromatic Leukodystrophy Methylmalonic Acidemia, cbIA Type	< 1 in 50,000 1 in 18,000 1 in 14,000 1 in 36,000 1 in 13,000 1 in 6,000 < 1 in 50,000 1 in 16,000 < 1 in 50,000	<pre>&lt; 1 in 1,000,000 &lt; 1 in 1,000,000</pre>
Leigh Syndrome, French-Canadian Type Lipoid Congenital Adrenal Hyperplasia Lysosomal Acid Lipase Deficiency Maple Syrup Urine Disease Type Ia Maple Syrup Urine Disease Type Ib Maple Syrup Urine Disease Type II Medium Chain Acyl-CoA Dehydrogenase Deficiency Megalencephalic Leukoencephalopathy with Subcortical Cysts Metachromatic Leukodystrophy Methylmalonic Acidemia, cbIA Type Methylmalonic Acidemia, cbIB Type	< 1 in 50,000 1 in 18,000 1 in 14,000 1 in 36,000 1 in 13,000 1 in 6,000 < 1 in 50,000 1 in 16,000 < 1 in 50,000 < 1 in 50,000 < 1 in 50,000 < 1 in 50,000	<pre>&lt; 1 in 1,000,000 &lt; 1 in 1,000,000</pre>
Leigh Syndrome, French-Canadian Type Lipoid Congenital Adrenal Hyperplasia Lysosomal Acid Lipase Deficiency Maple Syrup Urine Disease Type Ia Maple Syrup Urine Disease Type Ib Maple Syrup Urine Disease Type II Medium Chain Acyl-CoA Dehydrogenase Deficiency Megalencephalic Leukoencephalopathy with Subcortical Cysts Metachromatic Leukodystrophy Methylmalonic Acidemia, cbIA Type Methylmalonic Acidemia, cbIB Type Methylmalonic Aciduria and Homocystinuria, cbIC Type	< 1 in 50,000 1 in 18,000 1 in 14,000 1 in 36,000 1 in 13,000 1 in 6,000 < 1 in 50,000 1 in 16,000 < 1 in 50,000 < 1 in 50,000 < 1 in 50,000 1 in 16,000 < 1 in 50,000 1 in 16,000	< 1 in 1,000,000
Leigh Syndrome, French-Canadian Type Lipoid Congenital Adrenal Hyperplasia Lysosomal Acid Lipase Deficiency Maple Syrup Urine Disease Type Ia Maple Syrup Urine Disease Type Ib Maple Syrup Urine Disease Type II Medium Chain Acyl-CoA Dehydrogenase Deficiency Megalencephalic Leukoencephalopathy with Subcortical Cysts Metachromatic Leukodystrophy Methylmalonic Acidemia, cbIA Type Methylmalonic Acidemia, cbIB Type	< 1 in 50,000 1 in 18,000 1 in 14,000 1 in 36,000 1 in 13,000 1 in 6,000 < 1 in 50,000 1 in 16,000 < 1 in 50,000 < 1 in 50,000 < 1 in 50,000 < 1 in 50,000	<pre>&lt; 1 in 1,000,000 &lt; 1 in 1,000,000</pre>



MALE DONOR 10497

DOB: Ethnicity: Hispanic

Barcode: 11004512688138

FEMALE N/A

	DONOR 10497	
Disease	Residual Risk	Reproductive Risk
Mucolipidosis IV	< 1 in 50,000	< 1 in 1,000,000
Mucopolysaccharidosis Type I	1 in 16,000	< 1 in 1,000,000
Mucopolysaccharidosis Type II	< 1 in 1,000,000	1 in 300,000
Mucopolysaccharidosis Type IIIA	1 in 16,000	< 1 in 1,000,000
Mucopolysaccharidosis Type IIIB	1 in 31,000	< 1 in 1,000,000
Mucopolysaccharidosis Type IIIC	1 in 43,000	< 1 in 1,000,000
MUT-related Methylmalonic Acidemia	1 in 17,000	< 1 in 1,000,000
MYO7A-related Disorders	1 in 15,000	< 1 in 1,000,000
NEB-related Nemaline Myopathy	1 in 1,200	1 in 400,000
Nephrotic Syndrome, NPHS1-related	< 1 in 50,000	< 1 in 1,000,000
Nephrotic Syndrome, NPHS2-related	1 in 35,000	< 1 in 1,000,000
Niemann-Pick Disease Type C1	1 in 17,000	< 1 in 1,000,000
Niemann-Pick Disease Type C2	< 1 in 50,000	< 1 in 1,000,000
Niemann-Pick Disease, SMPD1-related	1 in 25,000	< 1 in 1,000,000
Nijmegen Breakage Syndrome	< 1 in 50,000	< 1 in 1,000,000
Ornithine Transcarbamylase Deficiency	< 1 in 1,000,000	1 in 140,000
PCCA-related Propionic Acidemia	1 in 4,200	< 1 in 1,000,000
PCCB-related Propionic Acidemia	1 in 22,000	< 1 in 1,000,000
PCDH15-related Disorders	1 in 3,300	< 1 in 1,000,000
Pendred Syndrome	1 in 6,400	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 1	1 in 16,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 3	1 in 44,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 4	1 in 9,300	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 5	< 1 in 71,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 6	< 1 in 50,000	< 1 in 1,000,000
Phenylalanine Hydroxylase Deficiency	1 in 7,100	< 1 in 1,000,000
POMGNT-related Disorders	< 1 in 12,000	< 1 in 1,000,000
Pompe Disease	1 in 6,500	< 1 in 1,000,000
PPT1-related Neuronal Ceroid Lipofuscinosis	1 in 7,700	< 1 in 1,000,000
Primary Carnitine Deficiency	1 in 16,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 1	1 in 13,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 2	< 1 in 50,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 3	1 in 20,000	< 1 in 1,000,000
Pycnodysostosis	1 in 43,000	< 1 in 1,000,000
Pyruvate Carboxylase Deficiency	1 in 25,000	< 1 in 1,000,000
Rhizomelic Chondrodysplasia Punctata Type 1	1 in 16,000	< 1 in 1,000,000
RTEL1-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Sandhoff Disease	1 in 30,000	< 1 in 1,000,000
Short-chain Acyl-CoA Dehydrogenase Deficiency	1 in 9,700	< 1 in 1,000,000
Sjogren-Larsson Syndrome	< 1 in 12,000	< 1 in 1,000,000
SLC26A2-related Disorders	1 in 14,000	< 1 in 1,000,000
Smith-Lemli-Opitz Syndrome	1 in 17,000	< 1 in 1,000,000
Spastic Paraplegia Type 15	< 1 in 50,000	< 1 in 1,000,000
	Negative for g.27134T>G SNP	
Spinal Muscular Atrophy	SMN1: 2 copies	1 in 820,000
	1 in 1,800	
Spondylothoracic Dysostosis	< 1 in 50,000	< 1 in 1,000,000
TGM1-related Autosomal Recessive Congenital Ichthyosis	1 in 22,000	< 1 in 1,000,000
TPP1-related Neuronal Ceroid Lipofuscinosis	1 in 30,000	< 1 in 1,000,000
Tyrosine Hydroxylase Deficiency	< 1 in 50,000	< 1 in 1,000,000
Tyrosinemia Type I	1 in 16,000	< 1 in 1,000,000
Tyrosinemia Type II	1 in 25,000	< 1 in 1,000,000
USH1C-related Disorders	1 in 35,000	< 1 in 1,000,000
USH2A-related Disorders	1 in 2,200	< 1 in 1,000,000
Usher Syndrome Type 3	1 in 41,000	< 1 in 1,000,000
Very-long-chain Acyl-CoA Dehydrogenase Deficiency	1 in 14,000	< 1 in 1,000,000
Wilson Disease	1 in 9,000	< 1 in 1,000,000
X-linked Adrenoleukodystrophy	1 in 90,000	1 in 42,000
X-linked Alport Syndrome	Not calculated	Not calculated
X-linked Congenital Adrenal Hypoplasia	< 1 in 1,000,000	< 1 in 1,000,000
X-linked Juvenile Retinoschisis	< 1 in 1,000,000	1 in 50,000
X-linked Myotubular Myopathy	Not calculated	Not calculated



MALE

DONOR 10497

DOB

Ethnicity: Hispanic Barcode: 11004512688138 FEMALE N/A

al Risk Reproductive Risk
00,000 1 in 200,000
000 < 1 in 1,000,000
00 < 1 in 1,000,000
)