



RESULTS RECIPIENT  
**SEATTLE SPERM BANK**  
Attn: Jeffrey Olliffe  
4915 25th Ave NE Ste 204w  
Seattle, WA 98105-5668  
Phone: (206) 588-1484  
Fax: (206) 466-4696  
NPI: 1306838271  
Report Date: 12/08/2019

MALE  
**DONOR 12462**  
DOB: [REDACTED]  
Ethnicity: Northern European  
Sample Type: EDTA Blood  
Date of Collection: 11/26/2019  
Date Received: 11/29/2019  
Date Tested: 12/08/2019  
Barcode: 11004512586951  
Accession ID: CSLNJMUJ2AZZYDC  
Indication: Egg or sperm donor

FEMALE  
N/A

# Foresight® Carrier Screen

**POSITIVE: CARRIER**

## ABOUT THIS TEST

The **Myriad Foresight Carrier Screen** utilizes sequencing, maximizing coverage across all DNA regions tested, to help you learn about your chance to have a child with a genetic disease.

## RESULTS SUMMARY

Risk Details	DONOR 12462	Partner
Panel Information	Foresight Carrier Screen Universal Panel Fundamental Plus Panel Fundamental Panel (175 conditions tested)	N/A
<b>POSITIVE: CARRIER</b> <b>Wilson Disease</b>  Reproductive Risk: 1 in 350 Inheritance: Autosomal Recessive	<b>CARRIER*</b> NM_000053.3(ATP7B):c. 3207C>A(H1069Q) heterozygote	The reproductive risk presented is based on a hypothetical pairing with a partner of the same ethnic group. Carrier testing should be considered. See "Next Steps".

\*Carriers generally do not experience symptoms.

No disease-causing mutations were detected in any other gene tested. A complete list of all conditions tested can be found on page 6.

## CLINICAL NOTES

- None

## NEXT STEPS

- Carrier testing should be considered for the diseases specified above for the patient's partner, as both parents must be carriers before a child is at high risk of developing the disease.
- Genetic counseling is recommended and patients may wish to discuss any positive results with blood relatives, as there is an increased chance that they are also carriers.

## POSITIVE: CARRIER

# Wilson Disease

**Reproductive risk: 1 in 350**  
Risk before testing: 1 in 30,000

**Gene:** ATP7B | **Inheritance Pattern:** Autosomal Recessive

Patient	DONOR 12462	No partner tested
<b>Result</b>	⊕ Carrier	N/A
<b>Variant(s)</b>	NM_000053.3(ATP7B):c.3207C>A(H1069Q) heterozygote	N/A
<b>Methodology</b>	Sequencing with copy number analysis	N/A
<b>Interpretation</b>	This individual is a carrier of Wilson disease. Carriers generally do not experience symptoms.	N/A
<b>Detection rate</b>	>99%	N/A
<b>Exons tested</b>	NM_000053:1-21.	N/A

## What is Wilson Disease?

Wilson disease is an inherited disease that causes the body to retain too much copper. Copper deposits in the liver, brain, kidneys, and eyes eventually cause tissue damage and scarring that makes the affected organs stop working properly. If not diagnosed and treated early, the condition causes organ failure and death.

Symptoms typically first appear in childhood or early adolescence, but they can appear as early as age 3 and as late as age 70. The most common symptom is liver disease, which first appears as fatigue, abdominal pain, or jaundice. In some cases, it progresses quickly to liver or kidney failure, and will require a liver transplant.

Symptoms can also include neurological problems such as tremors, poor coordination, loss of fine motor skills, problems walking, muscle rigidity in the body or face, or difficulty swallowing. Some people with the condition also develop psychiatric problems including depression, poor impulse control, phobias, aggression, and compulsive behavior. Wilson disease may also interfere with memory and attention span.

Copper deposits also accumulate in the eyes, creating characteristic brown circles around the colored part of the eye. These circles do not interfere with vision.

Even with ongoing treatment to remove excess copper from the body, people with Wilson disease sometimes develop arthritis, heart problems, and endocrine disorders caused by copper accumulation.

## How common is Wilson Disease?

Worldwide, approximately 1 in 30,000 people have Wilson disease. It is most common in China, Japan, and Sardinia, where it may affect as many as 1 in 10,000 people.

## How is Wilson Disease treated?

Wilson disease requires lifelong, regular treatment to remove copper from the body. Most people with the condition take a medication called penicillamine (brand name: Cuprimine or Depen) several times a day by mouth, combined with vitamin B6. This traps and removes copper from the body through the urine. However, some people react to penicillamine with fever, rash, and other serious complications. These people may be treated with other oral medications such as trientine or high-dose zinc. For individuals who do not respond to medication or have severe side effects, liver transplant is a final treatment option.

With careful treatment prior to the first symptom's appearance, most symptoms can be prevented. If treatment begins after symptoms appear, these symptoms can often show marked improvement. Stopping treatment, however, will cause health problems to return.

People with Wilson disease should not use copper cooking utensils. They should avoid foods high in copper, such as liver, chocolate, mushrooms, nuts, and shellfish. If they live in an area with copper water pipes, they should drink distilled water.

## What is the prognosis for a person with Wilson Disease?

With proper treatment, Wilson disease can be managed for many years after diagnosis. Its effect on lifespan is unclear.

## Methods and Limitations

**DONOR 12462 [Foresight Carrier Screen]:** Sequencing with copy number analysis, spinal muscular atrophy, and analysis of homologous regions.

### Sequencing with copy number analysis

High-throughput sequencing and read depth-based copy number analysis are used to analyze the listed exons, as well as selected intergenic and intronic regions, of the genes in the Conditions Tested section of the report. The region of interest (ROI) of the test comprises these regions, in addition to the 20 intronic bases flanking each exon. In a minority of cases where genomic features (e.g., long homopolymers) compromise calling fidelity, the affected intronic bases are not included in the ROI. The ROI is sequenced to high coverage and the sequences are compared to standards and references of normal variation. More than 99% of all bases in the ROI are sequenced at greater than the minimum read depth. Mutations may not be detected in areas of lower sequence coverage. Small insertions and deletions may not be as accurately determined as single nucleotide variants. Genes that have closely related pseudogenes may be addressed by a different method. *CFTR* and *DMD* testing includes analysis for both large (exon-level) deletions and duplications with an average sensitivity of 99%, while other genes are only analyzed for large deletions with a sensitivity of >75%. However, the sensitivity may be higher for selected founder deletions. The breakpoints of copy number variants and exons affected are estimated from probe positions. Only exons known to be included in the copy number variant are provided in the name. In some cases, the copy number variant may be larger or smaller than indicated. If *GJB2* is tested, two large upstream deletions which overlap *GJB6* and affect the expression of *GJB2*, *del(GJB6-D13S1830)* and *del(GJB6-D13S1854)*, are also analyzed. Mosaicism or somatic variants present at low levels may not be detected. If detected, these may not be reported.

Detection rates are determined by using literature to estimate the fraction of disease alleles, weighted by frequency, that the methodology is unable to detect. Detection rates only account for analytical sensitivity and certain variants that have been previously described in the literature may not be reported if there is insufficient evidence for pathogenicity. Detection rates do not account for the disease-specific rates of de novo mutations.

All variants that are a recognized cause of the disease will be reported. In addition, variants that have not previously been established as a recognized cause of disease may be identified. In these cases, only variants classified as "likely" pathogenic are reported. Likely pathogenic variants are described elsewhere in the report as "likely to have a negative impact on gene function". Likely pathogenic variants are evaluated and classified by assessing the nature of the variant and reviewing reports of allele frequencies in cases and controls, functional studies, variant annotation and effect prediction, and segregation studies. Exon level duplications are assumed to be in tandem and are classified according to their predicted effect on the reading frame. Benign variants, variants of uncertain significance, and variants not directly associated with the intended disease phenotype are not reported. Curation summaries of reported variants are available upon request.

### Spinal muscular atrophy

Targeted copy number analysis is used to determine the copy number of exon 7 of the *SMN1* gene relative to other genes. Other mutations may interfere with this analysis. Some individuals with two copies of *SMN1* are carriers with two *SMN1* genes on one chromosome and a *SMN1* deletion on the other chromosome. This is more likely in individuals who have 2 copies of the *SMN1* gene and are positive for the g.27134T>G SNP, which affects the reported residual risk; Ashkenazi Jewish or Asian patients with this genotype have a high post-test likelihood of being carriers for SMA and are reported as carriers. The g.27134T>G SNP is only reported in individuals who have 2 copies of *SMN1*.

### Analysis of homologous regions

A combination of high-throughput sequencing, read depth-based copy number analysis, and targeted genotyping is used to determine the number of functional gene copies and/or the presence of selected loss of function mutations in certain genes that have homology to other regions. The precise breakpoints of large deletions in these genes cannot be determined, but are estimated from copy number analysis. High numbers of pseudogene copies may interfere with this analysis.

If *CYP21A2* is tested, patients who have one or more additional copies of the *CYP21A2* gene and a loss of function mutation may not actually be a carrier of 21-hydroxylase-deficient congenital adrenal hyperplasia (CAH). Because the true incidence of non-classic CAH is unknown, the residual carrier and reproductive risk numbers on the report are only based on published incidences for classic CAH. However, the published prevalence of non-classic CAH is highest in individuals of Ashkenazi Jewish, Hispanic, Italian, and Yugoslav descent. Therefore, the residual and reproductive risks are likely an underestimate of overall chances for 21-hydroxylase-deficient CAH, especially in the aforementioned populations, as they do not account for non-classic CAH. If *HBA1/HBA2* are tested, some individuals with four alpha globin genes may be carriers, with three genes on one chromosome and a deletion on the other chromosome. This and similar, but rare, carrier states, where complementary changes exist in both the gene and a pseudogene, may not be detected by the assay.

## Limitations

In an unknown number of cases, nearby genetic variants may interfere with mutation detection. Other possible sources of diagnostic error include sample mix-up, trace contamination, bone marrow transplantation, blood transfusions and technical errors. This test is designed to detect and report germline alterations. While somatic variants present at low levels may be detected, these may not be reported. If more than one variant is detected in a gene, additional studies may be necessary to determine if those variants lie on the same chromosome or different chromosomes. The test does not fully address all inherited forms of intellectual disability, birth defects and genetic disease. A family history of any of these conditions may warrant additional evaluation. Furthermore, not all mutations will be identified in the genes analyzed and additional testing may be beneficial for some patients. For example, individuals of African, Southeast Asian, and Mediterranean ancestry are at increased risk for being carriers for hemoglobinopathies, which can be identified by CBC and hemoglobin electrophoresis or HPLC (*ACOG Practice Bulletin No. 78. Obstet. Gynecol. 2007;109:229-37*).

This test was developed and its performance characteristics determined by Myriad Women's Health, Inc. It has not been cleared or approved by the US Food and Drug Administration (FDA). The FDA does not require this test to go through premarket review. This test is used for clinical purposes. It should not be regarded as investigational or for research. This laboratory is certified under the Clinical Laboratory Improvement Amendments of 1988 (CLIA) as qualified to perform high-complexity clinical testing. These results are adjunctive to the ordering physician's evaluation. CLIA Number: **#05D1102604**.

## Resources

**GENOME CONNECT** | <http://www.genomeconnect.org>

Patients can share their reports via research registries such as Genome Connect, an online research registry working to build the knowledge base about genetics and health. Genome Connect provides patients, physicians, and researchers an opportunity to share genetic information to support the study of the impact of genetic variation on health conditions.

---

### SENIOR LABORATORY DIRECTOR



Jack Ji, PhD, FACMG

Report content approved by Jack Ji, PhD, FACMG on Dec 9, 2019

# Conditions Tested

## 11-beta-hydroxylase-deficient Congenital Adrenal Hyperplasia - Gene: CYP11B1.

Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000497:1-9. **Detection Rate:** Northern European 94%.

**6-pyruvoyl-tetrahydropterin Synthase Deficiency** - Gene: PTS. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000317:1-6. **Detection Rate:** Northern European >99%.

**ABCC8-related Familial Hyperinsulinism** - Gene: ABCC8. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000352:1-39. **Detection Rate:** Northern European >99%.

**Adenosine Deaminase Deficiency** - Gene: ADA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000022:1-12. **Detection Rate:** Northern European >99%.

**Alpha Thalassemia** - Genes: HBA1, HBA2. Autosomal Recessive. Analysis of homologous regions. **Variants (13):** -(alpha)20.5, --BRIT, --MEDI, --MEDII, --SEA, --THAI or --FIL, -alpha3.7, -alpha4.2, HBA1+HBA2 deletion, Hb Constant Spring, anti3.7, anti4.2, del HS-40. **Detection Rate:** Unknown due to rarity of disease.

**Alpha-mannosidosis** - Gene: MAN2B1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000528:1-23. **Detection Rate:** Northern European >99%.

**Alpha-sarcoglycanopathy** - Gene: SGCA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000023:1-9. **Detection Rate:** Northern European >99%.

**Alstrom Syndrome** - Gene: ALMS1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_015120:1-23. **Detection Rate:** Northern European >99%.

**AMT-related Glycine Encephalopathy** - Gene: AMT. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000481:1-9. **Detection Rate:** Northern European >99%.

**Andermann Syndrome** - Gene: SLC12A6. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_133647:1-25. **Detection Rate:** Northern European >99%.

**Argininemia** - Gene: ARG1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000045:1-8. **Detection Rate:** Northern European 97%.

**Argininosuccinic Aciduria** - Gene: ASL. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_001024943:1-16. **Detection Rate:** Northern European >99%.

**Aspartylglucosaminuria** - Gene: AGA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000027:1-9. **Detection Rate:** Northern European >99%.

**Ataxia with Vitamin E Deficiency** - Gene: TTPA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000370:1-5. **Detection Rate:** Northern European >99%.

**Ataxia-telangiectasia** - Gene: ATM. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000051:2-63. **Detection Rate:** Northern European 98%.

**ATP7A-related Disorders** - Gene: ATP7A. X-linked Recessive. Sequencing with copy number analysis. Exons: NM\_000052:2-23. **Detection Rate:** Northern European 96%.

**Autoimmune Polyglandular Syndrome Type 1** - Gene: AIRE. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000383:1-14. **Detection Rate:** Northern European >99%.

**Autosomal Recessive Osteopetrosis Type 1** - Gene: TCIRG1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_006019:2-20. **Detection Rate:** Northern European >99%.

**Autosomal Recessive Polycystic Kidney Disease, PKHD1-related** - Gene: PKHD1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_138694:2-67. **Detection Rate:** Northern European >99%.

## Autosomal Recessive Spastic Ataxia of Charlevoix-Saguenay - Gene: SACS.

Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_014363:2-10. **Detection Rate:** Northern European 99%.

**Bardet-Biedl Syndrome, BBS1-related** - Gene: BBS1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_024649:1-17. **Detection Rate:** Northern European >99%.

**Bardet-Biedl Syndrome, BBS10-related** - Gene: BBS10. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_024685:1-2. **Detection Rate:** Northern European >99%.

**Bardet-Biedl Syndrome, BBS12-related** - Gene: BBS12. Autosomal Recessive. Sequencing with copy number analysis. Exon: NM\_152618:2. **Detection Rate:** Northern European >99%.

**Bardet-Biedl Syndrome, BBS2-related** - Gene: BBS2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_031885:1-17. **Detection Rate:** Northern European >99%.

**BCS1L-related Disorders** - Gene: BCS1L. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_004328:3-9. **Detection Rate:** Northern European >99%.

**Beta-sarcoglycanopathy** - Gene: SGCB. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000232:1-6. **Detection Rate:** Northern European >99%.

**Biotinidase Deficiency** - Gene: BTD. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000060:1-4. **Detection Rate:** Northern European >99%.

**Bloom Syndrome** - Gene: BLM. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000057:2-22. **Detection Rate:** Northern European >99%.

**Calpainopathy** - Gene: CAPN3. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000070:1-24. **Detection Rate:** Northern European >99%.

**Canavan Disease** - Gene: ASPA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000049:1-6. **Detection Rate:** Northern European 98%.

**Carbamoylphosphate Synthetase I Deficiency** - Gene: CPS1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_001875:1-38. **Detection Rate:** Northern European >99%.

**Carnitine Palmitoyltransferase IA Deficiency** - Gene: CPT1A. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_001876:2-19. **Detection Rate:** Northern European >99%.

**Carnitine Palmitoyltransferase II Deficiency** - Gene: CPT2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000098:1-5. **Detection Rate:** Northern European >99%.

**Cartilage-hair Hypoplasia** - Gene: RMRP. Autosomal Recessive. Sequencing with copy number analysis. Exon: NR\_003051:1. **Detection Rate:** Northern European >99%.

**Cerebrotendinous Xanthomatosis** - Gene: CYP27A1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000784:1-9. **Detection Rate:** Northern European >99%.

**Citrullinemia Type 1** - Gene: ASS1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000050:3-16. **Detection Rate:** Northern European >99%.

**CLN3-related Neuronal Ceroid Lipofuscinosis** - Gene: CLN3. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_001042432:2-16. **Detection Rate:** Northern European >99%.

**CLN5-related Neuronal Ceroid Lipofuscinosis** - Gene: CLN5. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_006493:1-4. **Detection Rate:** Northern European >99%.

**CLN6-related Neuronal Ceroid Lipofuscinosis** - Gene: CLN6. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_017882:1-7. **Detection Rate:** Northern European >99%.

**CLN8-related Neuronal Ceroid Lipofuscinosis** - Gene: CLN8. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_018941:2-3. **Detection Rate:** Northern European >99%.

**Cohen Syndrome** - Gene: VPS13B. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_017890:2-62. **Detection Rate:** Northern European 97%.

**COL4A3-related Alport Syndrome** - Gene: COL4A3. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000091:1-52. **Detection Rate:** Northern European 97%.

**COL4A4-related Alport Syndrome** - Gene: COL4A4. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000092:2-48. **Detection Rate:** Northern European 98%.

**Combined Pituitary Hormone Deficiency, PROP1-related** - Gene: PROP1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_006261:1-3. **Detection Rate:** Northern European >99%.

**Congenital Adrenal Hyperplasia, CYP21A2-related** - Gene: CYP21A2. Autosomal Recessive. Analysis of homologous regions. **Variants (13):** CYP21A2 deletion, CYP21A2 duplication, CYP21A2 triplication, G111Vfs\*21, I173N, L308Ffs\*6, P31L, Q319\*, Q319\*+CYP21A2dup, R357W, V281L, [I237N;V238E;M240K], c.293-13C>G. **Detection Rate:** Northern European 96%.

**Congenital Disorder of Glycosylation Type Ia** - Gene: PMM2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000303:1-8. **Detection Rate:** Northern European >99%.

**Congenital Disorder of Glycosylation Type Ic** - Gene: ALG6. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_013339:2-15. **Detection Rate:** Northern European >99%.

**Congenital Disorder of Glycosylation, MPI-related** - Gene: MPI. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_002435:1-8. **Detection Rate:** Northern European >99%.

**Costeff Optic Atrophy Syndrome** - Gene: OPA3. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_025136:1-2. **Detection Rate:** Northern European >99%.

**Cystic Fibrosis** - Gene: CFTR. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000492:1-27. IVS8-5T allele analysis is only reported in the presence of the R117H mutation. **Detection Rate:** Northern European >99%.

**Cystinosis** - Gene: CTNS. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_004937:3-12. **Detection Rate:** Northern European >99%.

**D-bifunctional Protein Deficiency** - Gene: HSD17B4. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000414:1-24. **Detection Rate:** Northern European 98%.

**Delta-sarcoglycanopathy** - Gene: SGCD. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000337:2-9. **Detection Rate:** Northern European 99%.

**Dihydrolipoamide Dehydrogenase Deficiency** - Gene: DLD. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000108:1-14. **Detection Rate:** Northern European >99%.

**Dysferlinopathy** - Gene: DYSF. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_003494:1-55. **Detection Rate:** Northern European 98%.

**Dystrophinopathy (Including Duchenne/Becker Muscular Dystrophy)** - Gene: DMD. X-linked Recessive. Sequencing with copy number analysis. Exons: NM\_004006:1-79. **Detection Rate:** Northern European >99%.

**ERCC6-related Disorders** - Gene: ERCC6. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000124:2-21. **Detection Rate:** Northern European 99%.

**ERCC8-related Disorders** - Gene: ERCC8. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000082:1-12. **Detection Rate:** Northern European 95%.

**EVC-related Ellis-van Creveld Syndrome** - Gene: EVC. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_153717:1-21. **Detection Rate:** Northern European 96%.

**EVC2-related Ellis-van Creveld Syndrome** - Gene: EVC2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_147127:1-22. **Detection Rate:** Northern European >99%.

**Fabry Disease** - Gene: GLA. X-linked Recessive. Sequencing with copy number analysis. Exons: NM\_000169:1-7. **Detection Rate:** Northern European 98%.

**Familial Dysautonomia** - Gene: IKBKAP. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_003640:2-37. **Detection Rate:** Northern European >99%.

**Familial Mediterranean Fever** - Gene: MEFV. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000243:1-10. **Detection Rate:** Northern European >99%.

**Fanconi Anemia Complementation Group A** - Gene: FANCA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000135:1-43. **Detection Rate:** Northern European 92%.

**Fanconi Anemia, FANCC-related** - Gene: FANCC. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000136:2-15. **Detection Rate:** Northern European >99%.

**FKRP-related Disorders** - Gene: FKRP. Autosomal Recessive. Sequencing with copy number analysis. Exon: NM\_024301:4. **Detection Rate:** Northern European >99%.

**FKTN-related Disorders** - Gene: FKTN. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_001079802:3-11. **Detection Rate:** Northern European >99%.

**Galactokinase Deficiency** - Gene: GALK1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000154:1-8. **Detection Rate:** Northern European >99%.

**Galactosemia** - Gene: GALT. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000155:1-11. **Detection Rate:** Northern European >99%.

**Gamma-sarcoglycanopathy** - Gene: SGCG. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000231:2-8. **Detection Rate:** Northern European 88%.

**Gaucher Disease** - Gene: GBA. Autosomal Recessive. Analysis of homologous regions. **Variants (10):** D409V, D448H, IVS2+1G>A, L444P, N370S, R463C, R463H, R496H, V394L, p.L29Afs\*18. **Detection Rate:** Northern European 60%.

**GJB2-related DFNB1 Nonsyndromic Hearing Loss and Deafness** - Gene: GJB2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_004004:1-2. **Detection Rate:** Northern European >99%.

**GLB1-related Disorders** - Gene: GLB1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000404:1-16. **Detection Rate:** Northern European >99%.

**GLDC-related Glycine Encephalopathy** - Gene: GLDC. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000170:1-25. **Detection Rate:** Northern European 94%.

**Glutaric Acidemia, GCDH-related** - Gene: GCDH. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000159:2-12. **Detection Rate:** Northern European >99%.

**Glycogen Storage Disease Type Ia** - Gene: G6PC. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000151:1-5. **Detection Rate:** Northern European >99%.

**Glycogen Storage Disease Type Ib** - Gene: SLC37A4. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_001164277:3-11. **Detection Rate:** Northern European >99%.

**Glycogen Storage Disease Type III** - Gene: AGL. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000642:2-34. **Detection Rate:** Northern European >99%.

**GNE Myopathy** - Gene: GNE. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_001128227:1-12. **Detection Rate:** Northern European >99%.

**GNPTAB-related Disorders** - Gene: GNPTAB. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_024312:1-21. **Detection Rate:** Northern European >99%.

**HADHA-related Disorders** - Gene: HADHA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000182:1-20. **Detection Rate:** Northern European >99%.

**Hb Beta Chain-related Hemoglobinopathy (Including Beta Thalassemia and Sickle Cell Disease)** - Gene: HBB. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000518:1-3. **Detection Rate:** Northern European >99%.



**Hereditary Fructose Intolerance** - Gene: ALDOB. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000035:2-9. **Detection Rate:** Northern European >99%.

**Herlitz Junctional Epidermolysis Bullosa, LAMB3-related** - Gene: LAMB3. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000228:2-23. **Detection Rate:** Northern European >99%.

**Hexosaminidase A Deficiency (Including Tay-Sachs Disease)** - Gene: HEXA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000520:1-14. **Detection Rate:** Northern European >99%.

**HMG-CoA Lyase Deficiency** - Gene: HMGCL. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000191:1-9. **Detection Rate:** Northern European 98%.

**Holocarboxylase Synthetase Deficiency** - Gene: HLCS. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000411:4-12. **Detection Rate:** Northern European >99%.

**Homocystinuria, CBS-related** - Gene: CBS. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000071:3-17. **Detection Rate:** Northern European >99%.

**Hydroletharus Syndrome** - Gene: HYL1. Autosomal Recessive. Sequencing with copy number analysis. Exon: NM\_145014:4. **Detection Rate:** Northern European >99%.

**Hypophosphatasia** - Gene: ALPL. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000478:2-12. **Detection Rate:** Northern European >99%.

**Isovaleric Acidemia** - Gene: IVD. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_002225:1-12. **Detection Rate:** Northern European >99%.

**Joubert Syndrome 2** - Gene: TMEM216. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_001173990:1-5. **Detection Rate:** Northern European >99%.

**Junctional Epidermolysis Bullosa, LAMA3-related** - Gene: LAMA3. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000227:1-38. **Detection Rate:** Northern European >99%.

**Junctional Epidermolysis Bullosa, LAMC2-related** - Gene: LAMC2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_005562:1-23. **Detection Rate:** Northern European >99%.

**KCNJ11-related Familial Hyperinsulinism** - Gene: KCNJ11. Autosomal Recessive. Sequencing with copy number analysis. Exon: NM\_000525:1. **Detection Rate:** Northern European >99%.

**Krabbe Disease** - Gene: GALC. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000153:1-17. **Detection Rate:** Northern European >99%.

**LAMA2-related Muscular Dystrophy** - Gene: LAMA2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000426:1-65. **Detection Rate:** Northern European >99%.

**Leigh Syndrome, French-Canadian Type** - Gene: LRPPRC. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_133259:1-38. **Detection Rate:** Northern European >99%.

**Lipoid Congenital Adrenal Hyperplasia** - Gene: STAR. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000349:1-7. **Detection Rate:** Northern European >99%.

**Lysosomal Acid Lipase Deficiency** - Gene: LIPA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000235:2-10. **Detection Rate:** Northern European >99%.

**Maple Syrup Urine Disease Type Ia** - Gene: BCKDHA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000709:1-9. **Detection Rate:** Northern European >99%.

**Maple Syrup Urine Disease Type Ib** - Gene: BCKDHB. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_183050:1-10. **Detection Rate:** Northern European >99%.

**Maple Syrup Urine Disease Type II** - Gene: DBT. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_001918:1-11. **Detection Rate:** Northern European 96%.

**Medium Chain Acyl-CoA Dehydrogenase Deficiency** - Gene: ACADM. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000016:1-12. **Detection Rate:** Northern European >99%.

**Megalencephalic Leukoencephalopathy with Subcortical Cysts** - Gene: MLC1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_015166:2-12. **Detection Rate:** Northern European >99%.

**Metachromatic Leukodystrophy** - Gene: ARSA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000487:1-8. **Detection Rate:** Northern European >99%.

**Methylmalonic Acidemia, cblA Type** - Gene: MMAA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_172250:2-7. **Detection Rate:** Northern European >99%.

**Methylmalonic Acidemia, cblB Type** - Gene: MMAB. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_052845:1-9. **Detection Rate:** Northern European >99%.

**Methylmalonic Aciduria and Homocystinuria, cblC Type** - Gene: MMACHC. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_015506:1-4. **Detection Rate:** Northern European >99%.

**MKS1-related Disorders** - Gene: MKS1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_017777:1-18. **Detection Rate:** Northern European >99%.

**Mucopolidosis III Gamma** - Gene: GNPTG. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_032520:1-11. **Detection Rate:** Northern European >99%.

**Mucopolidosis IV** - Gene: MCOLN1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_020533:1-14. **Detection Rate:** Northern European >99%.

**Mucopolysaccharidosis Type I** - Gene: IDUA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000203:1-14. **Detection Rate:** Northern European >99%.

**Mucopolysaccharidosis Type II** - Gene: IDS. X-linked Recessive. Sequencing with copy number analysis. Exons: NM\_000202:1-9. **Detection Rate:** Northern European 88%.

**Mucopolysaccharidosis Type IIIA** - Gene: SGSH. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000199:1-8. **Detection Rate:** Northern European >99%.

**Mucopolysaccharidosis Type IIIB** - Gene: NAGLU. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000263:1-6. **Detection Rate:** Northern European >99%.

**Mucopolysaccharidosis Type IIIC** - Gene: HGSNAT. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_152419:1-18. **Detection Rate:** Northern European >99%.

**MUT-related Methylmalonic Acidemia** - Gene: MUT. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000255:2-13. **Detection Rate:** Northern European >99%.

**MYO7A-related Disorders** - Gene: MYO7A. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000260:2-49. **Detection Rate:** Northern European >99%.

**NEB-related Nemaline Myopathy** - Gene: NEB. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_001271208:3-80,117-183. **Detection Rate:** Northern European 92%.

**Nephrotic Syndrome, NPHS1-related** - Gene: NPHS1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_004646:1-29. **Detection Rate:** Northern European >99%.

**Nephrotic Syndrome, NPHS2-related** - Gene: NPHS2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_014625:1-8. **Detection Rate:** Northern European >99%.

**Niemann-Pick Disease Type C1** - Gene: NPC1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000271:1-25. **Detection Rate:** Northern European >99%.

**Niemann-Pick Disease Type C2** - Gene: NPC2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_006432:1-5. **Detection Rate:** Northern European >99%.



**Niemann-Pick Disease, SMPD1-related** - Gene: SMPD1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000543:1-6. **Detection Rate:** Northern European >99%.

**Nijmegen Breakage Syndrome** - Gene: NBN. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_002485:1-16. **Detection Rate:** Northern European >99%.

**Ornithine Transcarbamylase Deficiency** - Gene: OTC. X-linked Recessive. Sequencing with copy number analysis. Exons: NM\_000531:1-10. **Detection Rate:** Northern European 97%.

**PCCA-related Propionic Acidemia** - Gene: PCCA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000282:1-24. **Detection Rate:** Northern European 95%.

**PCCB-related Propionic Acidemia** - Gene: PCCB. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000532:1-15. **Detection Rate:** Northern European >99%.

**PCDH15-related Disorders** - Gene: PCDH15. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_033056:2-33. **Detection Rate:** Northern European 93%.

**Pendred Syndrome** - Gene: SLC26A4. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000441:2-21. **Detection Rate:** Northern European >99%.

**Peroxisome Biogenesis Disorder Type 1** - Gene: PEX1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000466:1-24. **Detection Rate:** Northern European >99%.

**Peroxisome Biogenesis Disorder Type 3** - Gene: PEX12. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000286:1-3. **Detection Rate:** Northern European >99%.

**Peroxisome Biogenesis Disorder Type 4** - Gene: PEX6. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000287:1-17. **Detection Rate:** Northern European 97%.

**Peroxisome Biogenesis Disorder Type 5** - Gene: PEX2. Autosomal Recessive. Sequencing with copy number analysis. Exon: NM\_000318:4. **Detection Rate:** Northern European >99%.

**Peroxisome Biogenesis Disorder Type 6** - Gene: PEX10. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_153818:1-6. **Detection Rate:** Northern European >99%.

**Phenylalanine Hydroxylase Deficiency** - Gene: PAH. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000277:1-13. **Detection Rate:** Northern European >99%.

**POMGNT-related Disorders** - Gene: POMGNT1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_017739:2-22. **Detection Rate:** Northern European 96%.

**Pompe Disease** - Gene: GAA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000152:2-20. **Detection Rate:** Northern European 98%.

**PPT1-related Neuronal Ceroid Lipofuscinosis** - Gene: PPT1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000310:1-9. **Detection Rate:** Northern European >99%.

**Primary Carnitine Deficiency** - Gene: SLC22A5. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_003060:1-10. **Detection Rate:** Northern European >99%.

**Primary Hyperoxaluria Type 1** - Gene: AGXT. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000030:1-11. **Detection Rate:** Northern European >99%.

**Primary Hyperoxaluria Type 2** - Gene: GRHPR. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_012203:1-9. **Detection Rate:** Northern European >99%.

**Primary Hyperoxaluria Type 3** - Gene: HOGA1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_138413:1-7. **Detection Rate:** Northern European >99%.

**Pycnodysostosis** - Gene: CTSK. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000396:2-8. **Detection Rate:** Northern European >99%.

**Pyruvate Carboxylase Deficiency** - Gene: PC. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000920:3-22. **Detection Rate:** Northern European >99%.

**Rhizomelic Chondrodysplasia Punctata Type 1** - Gene: PEX7. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000288:1-10. **Detection Rate:** Northern European >99%.

**RTEL1-related Disorders** - Gene: RTEL1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_032957:2-35. **Detection Rate:** Northern European >99%.

**Salla Disease** - Gene: SLC17A5. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_012434:1-11. **Detection Rate:** Northern European 98%.

**Sandhoff Disease** - Gene: HEXB. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000521:1-14. **Detection Rate:** Northern European >99%.

**Short-chain Acyl-CoA Dehydrogenase Deficiency** - Gene: ACADS. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000017:1-10. **Detection Rate:** Northern European >99%.

**Sjogren-Larsson Syndrome** - Gene: ALDH3A2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000382:1-10. **Detection Rate:** Northern European 96%.

**SLC26A2-related Disorders** - Gene: SLC26A2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000112:2-3. **Detection Rate:** Northern European >99%.

**Smith-Lemli-Opitz Syndrome** - Gene: DHCR7. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_001360:3-9. **Detection Rate:** Northern European >99%.

**Spastic Paraplegia Type 15** - Gene: ZFYVE26. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_015346:2-42. **Detection Rate:** Northern European >99%.

**Spinal Muscular Atrophy** - Gene: SMN1. Autosomal Recessive. Spinal muscular atrophy. Variant (1): SMN1 copy number. **Detection Rate:** Northern European 95%.

**Spondyl thoracic Dysostosis** - Gene: MESP2. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_001039958:1-2. **Detection Rate:** Northern European >99%.

**TGM1-related Autosomal Recessive Congenital Ichthyosis** - Gene: TGM1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000359:2-15. **Detection Rate:** Northern European >99%.

**TPP1-related Neuronal Ceroid Lipofuscinosis** - Gene: TPP1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000391:1-13. **Detection Rate:** Northern European >99%.

**Tyrosine Hydroxylase Deficiency** - Gene: TH. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_199292:1-14. **Detection Rate:** Northern European >99%.

**Tyrosinemia Type I** - Gene: FAH. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000137:1-14. **Detection Rate:** Northern European >99%.

**Tyrosinemia Type II** - Gene: TAT. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000353:2-12. **Detection Rate:** Northern European >99%.

**USH1C-related Disorders** - Gene: USH1C. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_005709:1-21. **Detection Rate:** Northern European >99%.

**USH2A-related Disorders** - Gene: USH2A. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_206933:2-72. **Detection Rate:** Northern European 94%.

**Usher Syndrome Type 3** - Gene: CLRN1. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_174878:1-3. **Detection Rate:** Northern European >99%.

**Very-long-chain Acyl-CoA Dehydrogenase Deficiency** - Gene: ACADVL. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000018:1-20. **Detection Rate:** Northern European >99%.



RESULTS RECIPIENT  
**SEATTLE SPERM BANK**  
Attn: Jeffrey Olliffe  
NPI: 1306838271  
Report Date: 12/08/2019

MALE  
**DONOR 12462**  
DOB: [REDACTED]  
Ethnicity: Northern European  
Barcode: 11004512586951

FEMALE  
N/A

**Wilson Disease** - Gene: ATP7B. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000053:1-21. **Detection Rate:** Northern European >99%.

**X-linked Adrenoleukodystrophy** - Gene: ABCD1. X-linked Recessive. Sequencing with copy number analysis. Exons: NM\_000033:1-6. **Detection Rate:** Northern European 77%.

**X-linked Alport Syndrome** - Gene: COL4A5. X-linked Recessive. Sequencing with copy number analysis. Exons: NM\_000495:1-51. **Detection Rate:** Northern European 95%.

**X-linked Congenital Adrenal Hypoplasia** - Gene: NR0B1. X-linked Recessive. Sequencing with copy number analysis. Exons: NM\_000475:1-2. **Detection Rate:** Northern European 99%.

**X-linked Juvenile Retinoschisis** - Gene: RS1. X-linked Recessive. Sequencing with copy number analysis. Exons: NM\_000330:1-6. **Detection Rate:** Northern European 98%.

**X-linked Myotubular Myopathy** - Gene: MTM1. X-linked Recessive. Sequencing with copy number analysis. Exons: NM\_000252:2-15. **Detection Rate:** Northern European 98%.

**X-linked Severe Combined Immunodeficiency** - Gene: IL2RG. X-linked Recessive. Sequencing with copy number analysis. Exons: NM\_000206:1-8. **Detection Rate:** Northern European >99%.

**Xeroderma Pigmentosum Group A** - Gene: XPA. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_000380:1-6. **Detection Rate:** Northern European >99%.

**Xeroderma Pigmentosum Group C** - Gene: XPC. Autosomal Recessive. Sequencing with copy number analysis. Exons: NM\_004628:1-16. **Detection Rate:** Northern European 97%.

## Risk Calculations

Below are the risk calculations for all conditions tested. Since negative results do not completely rule out the possibility of being a carrier, the **residual risk** represents the patient's post-test likelihood of being a carrier and the **reproductive risk** represents the likelihood the patient's future children could inherit each disease. These risks are inherent to all carrier screening tests, may vary by ethnicity, are predicated on a negative family history and are present even after a negative test result. Inaccurate reporting of ethnicity may cause errors in risk calculation. The reproductive risk presented is based on a hypothetical pairing with a partner of the same ethnic group.

†Indicates a positive result. See the full clinical report for interpretation and details.

Disease	DONOR 12462 Residual Risk	Reproductive Risk
11-beta-hydroxylase-deficient Congenital Adrenal Hyperplasia	1 in 3,800	< 1 in 1,000,000
6-pyruvoyl-tetrahydropterin Synthase Deficiency	< 1 in 50,000	< 1 in 1,000,000
ABCC8-related Familial Hyperinsulinism	1 in 17,000	< 1 in 1,000,000
Adenosine Deaminase Deficiency	1 in 22,000	< 1 in 1,000,000
Alpha Thalassemia	Alpha globin status: aa/aa.	Not calculated
Alpha-mannosidosis	1 in 35,000	< 1 in 1,000,000
Alpha-sarcoglycanopathy	1 in 45,000	< 1 in 1,000,000
Alstrom Syndrome	< 1 in 50,000	< 1 in 1,000,000
AMT-related Glycine Encephalopathy	1 in 22,000	< 1 in 1,000,000
Andermann Syndrome	< 1 in 50,000	< 1 in 1,000,000
Argininemia	< 1 in 17,000	< 1 in 1,000,000
Argininosuccinic Aciduria	1 in 13,000	< 1 in 1,000,000
Aspartylglucosaminuria	< 1 in 50,000	< 1 in 1,000,000
Ataxia with Vitamin E Deficiency	< 1 in 50,000	< 1 in 1,000,000
Ataxia-telangiectasia	1 in 11,000	< 1 in 1,000,000
ATP7A-related Disorders	< 1 in 1,000,000	1 in 600,000
Autoimmune Polyglandular Syndrome Type 1	1 in 15,000	< 1 in 1,000,000
Autosomal Recessive Osteopetrosis Type 1	1 in 35,000	< 1 in 1,000,000
Autosomal Recessive Polycystic Kidney Disease, PKHD1-related	1 in 8,100	< 1 in 1,000,000
Autosomal Recessive Spastic Ataxia of Charlevoix-Saguenay	< 1 in 44,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS1-related	1 in 32,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS10-related	1 in 42,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS12-related	< 1 in 50,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS2-related	< 1 in 50,000	< 1 in 1,000,000
BCS1L-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Beta-sarcoglycanopathy	1 in 39,000	< 1 in 1,000,000
Biotinidase Deficiency	1 in 13,000	1 in 650,000
Bloom Syndrome	< 1 in 50,000	< 1 in 1,000,000
Calpainopathy	1 in 13,000	< 1 in 1,000,000
Canavan Disease	1 in 9,700	< 1 in 1,000,000
Carbamoylphosphate Synthetase I Deficiency	< 1 in 57,000	< 1 in 1,000,000
Carnitine Palmitoyltransferase IA Deficiency	< 1 in 50,000	< 1 in 1,000,000
Carnitine Palmitoyltransferase II Deficiency	1 in 25,000	< 1 in 1,000,000
Cartilage-hair Hypoplasia	< 1 in 50,000	< 1 in 1,000,000
Cerebrotendinous Xanthomatosis	1 in 11,000	< 1 in 1,000,000
Citrullinemia Type 1	1 in 14,000	< 1 in 1,000,000
CLN3-related Neuronal Ceroid Lipofuscinosis	1 in 8,600	< 1 in 1,000,000
CLN5-related Neuronal Ceroid Lipofuscinosis	< 1 in 50,000	< 1 in 1,000,000
CLN6-related Neuronal Ceroid Lipofuscinosis	1 in 43,000	< 1 in 1,000,000
CLN8-related Neuronal Ceroid Lipofuscinosis	< 1 in 50,000	< 1 in 1,000,000
Cohen Syndrome	< 1 in 15,000	< 1 in 1,000,000
COL4A3-related Alport Syndrome	1 in 6,200	< 1 in 1,000,000
COL4A4-related Alport Syndrome	1 in 12,000	< 1 in 1,000,000
Combined Pituitary Hormone Deficiency, PROP1-related	1 in 6,100	< 1 in 1,000,000
Congenital Adrenal Hyperplasia, CYP21A2-related	1 in 1,300	1 in 280,000
Congenital Disorder of Glycosylation Type Ia	1 in 16,000	< 1 in 1,000,000
Congenital Disorder of Glycosylation Type Ic	< 1 in 50,000	< 1 in 1,000,000
Congenital Disorder of Glycosylation, MPI-related	< 1 in 50,000	< 1 in 1,000,000
Costeff Optic Atrophy Syndrome	< 1 in 50,000	< 1 in 1,000,000

Disease	DONOR 12462 Residual Risk	Reproductive Risk
Cystic Fibrosis	1 in 3,000	1 in 360,000
Cystinosis	1 in 22,000	< 1 in 1,000,000
D-bifunctional Protein Deficiency	1 in 9,000	< 1 in 1,000,000
Delta-sarcoglycanopathy	< 1 in 40,000	< 1 in 1,000,000
Dihydrolipoamide Dehydrogenase Deficiency	< 1 in 50,000	< 1 in 1,000,000
Dysferlinopathy	1 in 11,000	< 1 in 1,000,000
Dystrophinopathy (Including Duchenne/Becker Muscular Dystrophy)	Not calculated	Not calculated
ERCC6-related Disorders	1 in 26,000	< 1 in 1,000,000
ERCC8-related Disorders	< 1 in 9,900	< 1 in 1,000,000
EVC-related Ellis-van Creveld Syndrome	1 in 7,500	< 1 in 1,000,000
EVC2-related Ellis-van Creveld Syndrome	< 1 in 50,000	< 1 in 1,000,000
Fabry Disease	< 1 in 1,000,000	1 in 80,000
Familial Dysautonomia	< 1 in 50,000	< 1 in 1,000,000
Familial Mediterranean Fever	< 1 in 50,000	< 1 in 1,000,000
Fanconi Anemia Complementation Group A	1 in 2,800	< 1 in 1,000,000
Fanconi Anemia, FANCC-related	< 1 in 50,000	< 1 in 1,000,000
FKRP-related Disorders	1 in 16,000	< 1 in 1,000,000
FKTN-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Galactokinase Deficiency	1 in 10,000	< 1 in 1,000,000
Galactosemia	1 in 8,600	< 1 in 1,000,000
Gamma-sarcoglycanopathy	1 in 3,000	< 1 in 1,000,000
Gaucher Disease	1 in 260	1 in 110,000
GJB2-related DFNB1 Nonsyndromic Hearing Loss and Deafness	1 in 2,500	1 in 260,000
GLB1-related Disorders	1 in 19,000	< 1 in 1,000,000
GLDC-related Glycine Encephalopathy	1 in 2,800	< 1 in 1,000,000
Glutaric Acidemia, GCDH-related	1 in 16,000	< 1 in 1,000,000
Glycogen Storage Disease Type Ia	1 in 18,000	< 1 in 1,000,000
Glycogen Storage Disease Type Ib	1 in 35,000	< 1 in 1,000,000
Glycogen Storage Disease Type III	1 in 16,000	< 1 in 1,000,000
GNE Myopathy	1 in 23,000	< 1 in 1,000,000
GNPTAB-related Disorders	1 in 32,000	< 1 in 1,000,000
HADHA-related Disorders	1 in 20,000	< 1 in 1,000,000
Hb Beta Chain-related Hemoglobinopathy (Including Beta Thalassemia and Sickle Cell Disease)	1 in 3,100	1 in 390,000
Hereditary Fructose Intolerance	1 in 7,900	< 1 in 1,000,000
Herlitz Junctional Epidermolysis Bullosa, LAMB3-related	< 1 in 50,000	< 1 in 1,000,000
Hexosaminidase A Deficiency (Including Tay-Sachs Disease)	1 in 30,000	< 1 in 1,000,000
HMG-CoA Lyase Deficiency	< 1 in 33,000	< 1 in 1,000,000
Holocarboxylase Synthetase Deficiency	1 in 15,000	< 1 in 1,000,000
Homocystinuria, CBS-related	1 in 9,400	< 1 in 1,000,000
Hydrolethalus Syndrome	< 1 in 50,000	< 1 in 1,000,000
Hypophosphatasia	1 in 27,000	< 1 in 1,000,000
Isovaleric Acidemia	1 in 32,000	< 1 in 1,000,000
Joubert Syndrome 2	< 1 in 50,000	< 1 in 1,000,000
Junctional Epidermolysis Bullosa, LAMA3-related	< 1 in 50,000	< 1 in 1,000,000
Junctional Epidermolysis Bullosa, LAMC2-related	< 1 in 50,000	< 1 in 1,000,000
KCNJ11-related Familial Hyperinsulinism	< 1 in 50,000	< 1 in 1,000,000
Krabbe Disease	1 in 14,000	< 1 in 1,000,000
LAMA2-related Muscular Dystrophy	1 in 34,000	< 1 in 1,000,000
Leigh Syndrome, French-Canadian Type	< 1 in 50,000	< 1 in 1,000,000
Lipoid Congenital Adrenal Hyperplasia	< 1 in 50,000	< 1 in 1,000,000
Lysosomal Acid Lipase Deficiency	1 in 18,000	< 1 in 1,000,000
Maple Syrup Urine Disease Type Ia	1 in 42,000	< 1 in 1,000,000
Maple Syrup Urine Disease Type Ib	1 in 39,000	< 1 in 1,000,000
Maple Syrup Urine Disease Type II	1 in 13,000	< 1 in 1,000,000
Medium Chain Acyl-CoA Dehydrogenase Deficiency	1 in 4,400	1 in 790,000
Megalencephalic Leukoencephalopathy with Subcortical Cysts	< 1 in 50,000	< 1 in 1,000,000
Metachromatic Leukodystrophy	1 in 16,000	< 1 in 1,000,000
Methylmalonic Acidemia, cblA Type	< 1 in 50,000	< 1 in 1,000,000
Methylmalonic Acidemia, cblB Type	1 in 48,000	< 1 in 1,000,000
Methylmalonic Aciduria and Homocystinuria, cblC Type	1 in 16,000	< 1 in 1,000,000
MKS1-related Disorders	< 1 in 50,000	< 1 in 1,000,000

Disease	DONOR 12462 Residual Risk	Reproductive Risk
Mucopolipidosis III Gamma	< 1 in 50,000	< 1 in 1,000,000
Mucopolipidosis IV	< 1 in 50,000	< 1 in 1,000,000
Mucopolysaccharidosis Type I	1 in 16,000	< 1 in 1,000,000
Mucopolysaccharidosis Type II	1 in 600,000	1 in 150,000
Mucopolysaccharidosis Type IIIA	1 in 12,000	< 1 in 1,000,000
Mucopolysaccharidosis Type IIIB	1 in 25,000	< 1 in 1,000,000
Mucopolysaccharidosis Type IIIC	1 in 37,000	< 1 in 1,000,000
MUT-related Methylmalonic Acidemia	1 in 26,000	< 1 in 1,000,000
MYO7A-related Disorders	1 in 15,000	< 1 in 1,000,000
NEB-related Nemaline Myopathy	1 in 1,200	1 in 400,000
Nephrotic Syndrome, NPHS1-related	< 1 in 50,000	< 1 in 1,000,000
Nephrotic Syndrome, NPHS2-related	1 in 35,000	< 1 in 1,000,000
Niemann-Pick Disease Type C1	1 in 19,000	< 1 in 1,000,000
Niemann-Pick Disease Type C2	< 1 in 50,000	< 1 in 1,000,000
Niemann-Pick Disease, SMPD1-related	1 in 25,000	< 1 in 1,000,000
Nijmegen Breakage Syndrome	1 in 16,000	< 1 in 1,000,000
Ornithine Transcarbamylase Deficiency	< 1 in 1,000,000	1 in 140,000
PCCA-related Propionic Acidemia	1 in 4,200	< 1 in 1,000,000
PCCB-related Propionic Acidemia	1 in 22,000	< 1 in 1,000,000
PCDH15-related Disorders	1 in 3,300	< 1 in 1,000,000
Pendred Syndrome	1 in 8,200	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 1	1 in 16,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 3	1 in 44,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 4	1 in 9,300	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 5	< 1 in 71,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 6	< 1 in 50,000	< 1 in 1,000,000
Phenylalanine Hydroxylase Deficiency	1 in 4,800	1 in 940,000
POMGNT-related Disorders	< 1 in 12,000	< 1 in 1,000,000
Pompe Disease	1 in 4,000	< 1 in 1,000,000
PPT1-related Neuronal Ceroid Lipofuscinosis	1 in 7,700	< 1 in 1,000,000
Primary Carnitine Deficiency	1 in 11,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 1	1 in 17,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 2	< 1 in 50,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 3	1 in 13,000	< 1 in 1,000,000
Pycnodysostosis	1 in 43,000	< 1 in 1,000,000
Pyruvate Carboxylase Deficiency	1 in 25,000	< 1 in 1,000,000
Rhizomelic Chondrodysplasia Punctata Type 1	1 in 16,000	< 1 in 1,000,000
RTEL1-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Salla Disease	< 1 in 30,000	< 1 in 1,000,000
Sandhoff Disease	1 in 32,000	< 1 in 1,000,000
Short-chain Acyl-CoA Dehydrogenase Deficiency	1 in 11,000	< 1 in 1,000,000
Sjogren-Larsson Syndrome	< 1 in 12,000	< 1 in 1,000,000
SLC26A2-related Disorders	1 in 16,000	< 1 in 1,000,000
Smith-Lemli-Opitz Syndrome	1 in 9,400	< 1 in 1,000,000
Spastic Paraplegia Type 15	< 1 in 50,000	< 1 in 1,000,000
Spinal Muscular Atrophy	Negative for g.27134T>G SNP SMN1: 2 copies 1 in 770	1 in 110,000
Spondylothoracic Dysostosis	< 1 in 50,000	< 1 in 1,000,000
TGM1-related Autosomal Recessive Congenital Ichthyosis	1 in 22,000	< 1 in 1,000,000
TPP1-related Neuronal Ceroid Lipofuscinosis	1 in 30,000	< 1 in 1,000,000
Tyrosine Hydroxylase Deficiency	< 1 in 50,000	< 1 in 1,000,000
Tyrosinemia Type I	1 in 16,000	< 1 in 1,000,000
Tyrosinemia Type II	1 in 25,000	< 1 in 1,000,000
USH1C-related Disorders	1 in 35,000	< 1 in 1,000,000
USH2A-related Disorders	1 in 2,200	< 1 in 1,000,000
Usher Syndrome Type 3	1 in 41,000	< 1 in 1,000,000
Very-long-chain Acyl-CoA Dehydrogenase Deficiency	1 in 18,000	< 1 in 1,000,000
Wilson Disease	H1069Q heterozygote †	1 in 350
X-linked Adrenoleukodystrophy	1 in 90,000	1 in 42,000
X-linked Alport Syndrome	Not calculated	Not calculated
X-linked Congenital Adrenal Hypoplasia	< 1 in 1,000,000	< 1 in 1,000,000



RESULTS RECIPIENT  
SEATTLE SPERM BANK  
Attn: Jeffrey Olliffe  
NPI: 1306838271  
Report Date: 12/08/2019

MALE  
DONOR 12462  
DOB: [REDACTED]  
Ethnicity: Northern European  
Barcode: 11004512586951

FEMALE  
N/A

Disease	DONOR 12462 Residual Risk	Reproductive Risk
X-linked Juvenile Retinoschisis	< 1 in 1,000,000	1 in 40,000
X-linked Myotubular Myopathy	Not calculated	Not calculated
X-linked Severe Combined Immunodeficiency	< 1 in 1,000,000	1 in 200,000
Xeroderma Pigmentosum Group A	< 1 in 50,000	< 1 in 1,000,000
Xeroderma Pigmentosum Group C	1 in 7,300	< 1 in 1,000,000